

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 50 mg hard capsules
Tasigna 200 mg hard capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Tasigna 50 mg hard capsules

One hard capsule contains 50 mg nilotinib (as hydrochloride monohydrate).

Excipient with known effect

One hard capsule contains 39.03 mg lactose monohydrate.

Tasigna 200 mg hard capsules

One hard capsule contains 200 mg nilotinib (as hydrochloride monohydrate).

Excipient with known effect

One hard capsule contains 156.11 mg lactose monohydrate.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule.

Tasigna 50 mg hard capsules

White to yellowish powder in hard gelatin capsule with red opaque cap and light yellow opaque body, size 4 with black radial imprint "NVR/ABL" on cap.

Tasigna 200 mg hard capsules

White to yellowish powder in light yellow opaque hard gelatin capsules, size 0 with red axial imprint "NVR/TKI".

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Tasigna is indicated for the treatment of:

- adult and paediatric patients with newly diagnosed Philadelphia chromosome positive chronic myelogenous leukaemia (CML) in the chronic phase,
- adult patients with chronic phase and accelerated phase Philadelphia chromosome positive CML with resistance or intolerance to prior therapy including imatinib. Efficacy data in patients with CML in blast crisis are not available,
- paediatric patients with chronic phase Philadelphia chromosome positive CML with resistance or intolerance to prior therapy including imatinib.

4.2 Posology and method of administration

Therapy should be initiated by a physician experienced in the diagnosis and the treatment of patients with CML.

Posology

Treatment should be continued as long as clinical benefit is observed or until unacceptable toxicity occurs.

If a dose is missed the patient should not take an additional dose, but take the usual prescribed next dose.

Posology for Philadelphia chromosome positive CML adult patients

The recommended dose is:

- 300 mg twice daily in newly diagnosed patients with CML in the chronic phase,
- 400 mg twice daily in patients with chronic or accelerated phase CML with resistance or intolerance to prior therapy.

For a dose of 300 mg twice daily, 150 mg hard capsules are available.

Posology for Philadelphia chromosome positive CML paediatric patients

Dosing in paediatric patients is individualised and is based on body surface area (mg/m²). The recommended dose of nilotinib is 230 mg/m² twice daily, rounded to the nearest 50 mg dose (to a maximum single dose of 400 mg) (see Table 1). Different strengths of Tasigna hard capsules can be combined to attain the desired dose.

There is no experience with treatment of paediatric patients below 2 years of age. There are no data in newly diagnosed paediatric patients below 10 years of age and limited data in imatinib-resistant or intolerant paediatric patients below 6 years of age.

Table 1 Dosing scheme of nilotinib 230 mg/m² twice daily

Body Surface Area (BSA)	Dose in mg (twice daily)
Up to 0.32 m ²	50 mg
0.33 – 0.54 m ²	100 mg
0.55 – 0.76 m ²	150 mg
0.77 – 0.97 m ²	200 mg
0.98 – 1.19 m ²	250 mg
1.20 – 1.41 m ²	300 mg
1.42 – 1.63 m ²	350 mg
≥1.64 m ²	400 mg

Philadelphia chromosome positive CML patients in chronic phase who have been treated with nilotinib as first-line therapy and who achieved a sustained deep molecular response (MR4.5)

Discontinuation of treatment may be considered in eligible Philadelphia chromosome positive (Ph+) CML patients in chronic phase who have been treated with nilotinib at 300 mg twice daily for a minimum of 3 years if a deep molecular response is sustained for a minimum of one year immediately prior to discontinuation of therapy. Discontinuation of nilotinib therapy should be initiated by a physician experienced in the treatment of patients with CML (see sections 4.4 and 5.1).

Eligible patients who discontinue nilotinib therapy must have their BCR-ABL transcript levels and complete blood count with differential monitored monthly for one year, then every 6 weeks for the second year, and every 12 weeks thereafter. Monitoring of BCR-ABL transcript levels must be performed with a quantitative diagnostic test validated to measure molecular response levels on the International Scale (IS) with a sensitivity of at least MR4.5 (BCR-ABL/ABL ≤0.0032% IS).

For patients who lose MR4 (MR4=BCR-ABL/ABL ≤0.01%IS) but not MMR (MMR=BCR-ABL/ABL ≤0.1%IS) during the treatment-free phase, BCR-ABL transcript levels should be monitored every 2 weeks until BCR-ABL levels return to a range between MR4 and MR4.5.

Patients who maintain BCR-ABL levels between MMR and MR4 for a minimum of 4 consecutive measurements can return to the original monitoring schedule.

Patients who lose MMR must re-initiate treatment within 4 weeks of when loss of remission is known to have occurred. Nilotinib therapy should be re-initiated at 300 mg twice daily or at a reduced dose level of 400 mg once daily if the patient had a dose reduction prior to discontinuation of therapy. Patients who re-initiate nilotinib therapy should have their BCR-ABL transcript levels monitored monthly until MMR is re-established and every 12 weeks thereafter (see section 4.4).

Philadelphia chromosome positive CML patients in chronic phase who have achieved a sustained deep molecular response (MR 4.5) on nilotinib following prior imatinib therapy

Discontinuation of treatment may be considered in eligible Philadelphia chromosome positive (Ph+) CML patients in chronic phase who have been treated with nilotinib for a minimum of 3 years if a deep molecular response is sustained for a minimum of one year immediately prior to discontinuation of therapy. Discontinuation of nilotinib therapy should be initiated by a physician experienced in the treatment of patients with CML (see sections 4.4 and 5.1).

Eligible patients who discontinue nilotinib therapy must have their BCR-ABL transcript levels and complete blood count with differential monitored monthly for one year, then every 6 weeks for the second year, and every 12 weeks thereafter. Monitoring of BCR-ABL transcript levels must be performed with a quantitative diagnostic test validated to measure molecular response levels on the International Scale (IS) with a sensitivity of at least MR4.5 ($\text{BCR-ABL/ABL} \leq 0.0032\% \text{ IS}$).

Patients with confirmed loss of MR4 ($\text{MR4} = \text{BCR-ABL/ABL} \leq 0.01\% \text{ IS}$) during the treatment-free phase (two consecutive measures separated by at least 4 weeks showing loss of MR4) or loss of major molecular response ($\text{MMR} = \text{BCR-ABL/ABL} \leq 0.1\% \text{ IS}$) must re-initiate treatment within 4 weeks of when loss of remission is known to have occurred. Nilotinib therapy should be re-initiated at either 300 mg or 400 mg twice daily. Patients who re-initiate nilotinib therapy should have their BCR-ABL transcript levels monitored monthly until previous major molecular response or MR4 level is re-established and every 12 weeks thereafter (see section 4.4).

Dose adjustments or modifications

Tasigna may need to be temporarily withheld and/or dose reduced for haematological toxicities (neutropenia, thrombocytopenia) that are not related to the underlying leukaemia (see Table 2).

Table 2 Dose adjustments for neutropenia and thrombocytopenia

Adult patients with newly diagnosed chronic phase CML at 300 mg twice daily and imatinib-resistant or intolerant CML in chronic phase at 400 mg twice daily	ANC* <1.0 x 10 ⁹ /l and/or platelet counts <50 x 10 ⁹ /l	<ol style="list-style-type: none"> 1. Treatment with nilotinib must be interrupted and blood count monitored. 2. Treatment must be resumed within 2 weeks at prior dose if ANC >1.0 x 10⁹/l and/or platelets >50 x 10⁹/l. 3. If blood counts remain low, a dose reduction to 400 mg once daily may be required.
Adult patients with imatinib-resistant or intolerant CML in accelerated phase at 400 mg twice daily	ANC* <0.5 x 10 ⁹ /l and/or platelet counts <10 x 10 ⁹ /l	<ol style="list-style-type: none"> 1. Treatment with nilotinib must be interrupted and blood count monitored. 2. Treatment must be resumed within 2 weeks at prior dose if ANC >1.0 x 10⁹/l and/or platelets >20 x 10⁹/l. 3. If blood counts remain low, a dose reduction to 400 mg once daily may be required.
Paediatric patients with newly-diagnosed CML in chronic phase at 230 mg/m ² twice daily and imatinib-resistant or intolerant CML in chronic phase at 230 mg/m ² twice daily	ANC* <1.0 x 10 ⁹ /l and/or platelet counts <50 x 10 ⁹ /l	<ol style="list-style-type: none"> 1. Treatment with nilotinib must be interrupted and blood count monitored. 2. Treatment must be resumed within 2 weeks at prior dose if ANC >1.5 x 10⁹/l and/or platelets >75 x 10⁹/l. 3. If blood counts remain low, a dose reduction to 230 mg/m² once daily may be required. 4. If event occurs after dose reduction, consider discontinuing treatment.

*ANC = absolute neutrophil count

If clinically significant moderate or severe non-haematological toxicity develops, dosing should be interrupted, and patients should be monitored and treated accordingly. If the prior dose was 300 mg twice daily in adult newly diagnosed patients with CML in the chronic phase, or 400 mg twice daily in adult patients with imatinib-resistant or intolerant CML in chronic or accelerated phase, or 230 mg/m² twice daily in paediatric patients, dosing may be resumed at 400 mg once daily in adult patients and at 230 mg/m² once daily in paediatric patients once the toxicity has resolved. If the prior dose was 400 mg once daily in adult patients or 230 mg/m² once daily in paediatric patients, treatment should be discontinued. If clinically appropriate, re-escalation of the dose to the starting dose of 300 mg twice daily in adult newly diagnosed patients with CML in the chronic phase or to 400 mg twice daily in adult patients with imatinib-resistant or intolerant CML in chronic or accelerated phase or to 230 mg/m² twice daily in paediatric patients should be considered.

Elevated serum lipase: For Grade 3-4 serum lipase elevations, doses in adult patients should be reduced to 400 mg once daily or interrupted. In paediatric patients, treatment must be interrupted until the event returns to Grade ≤1. Thereafter, if the prior dose was 230 mg/m² twice daily, treatment can be resumed at 230 mg/m² once daily. If the prior dose was 230 mg/m² once daily, treatment should be discontinued. Serum lipase levels should be tested monthly or as clinically indicated (see section 4.4).

Elevated bilirubin and hepatic transaminases: For Grade 3-4 bilirubin and hepatic transaminase elevations in adult patients, doses should be reduced to 400 mg once daily or interrupted. For Grade ≥2 bilirubin elevations or Grade ≥3 hepatic transaminase elevations in paediatric patients, treatment must be interrupted until the levels return to Grade ≤1. Thereafter, if the prior dose was 230 mg/m² twice daily, treatment can be resumed at 230 mg/m² once daily. If the prior dose was 230 mg/m² once daily, and recovery to Grade ≤1 takes longer than 28 days, treatment should be discontinued. Bilirubin and hepatic transaminases levels should be tested monthly or as clinically indicated.

Special populations

Elderly

Approximately 12% of subjects in the Phase III study in patients with newly diagnosed CML in chronic phase and approximately 30% of subjects in the Phase II study in patients with imatinib-resistant or intolerant CML in chronic phase and accelerated phase were 65 years of age or over. No major differences were observed for safety and efficacy in patients ≥ 65 years of age as compared to adults aged 18 to 65 years.

Renal impairment

Clinical studies have not been performed in patients with impaired renal function. Since nilotinib and its metabolites are not renally excreted, a decrease in total body clearance is not anticipated in patients with renal impairment.

Hepatic impairment

Hepatic impairment has a modest effect on the pharmacokinetics of nilotinib. Dose adjustment is not considered necessary in patients with hepatic impairment. However, patients with hepatic impairment should be treated with caution (see section 4.4).

Cardiac disorders

In clinical studies, patients with uncontrolled or significant cardiac disease (e.g. recent myocardial infarction, congestive heart failure, unstable angina or clinically significant bradycardia) were excluded. Caution should be exercised in patients with relevant cardiac disorders (see section 4.4).

Increases in total serum cholesterol levels have been reported with nilotinib therapy (see section 4.4). Lipid profiles should be determined prior to initiating nilotinib therapy, assessed at month 3 and 6 after initiating therapy and at least yearly during chronic therapy.

Increases in blood glucose levels have been reported with nilotinib therapy (see section 4.4). Blood glucose levels should be assessed prior to initiating nilotinib therapy and monitored during treatment.

Paediatric population

The safety and efficacy of Tasigna in paediatric patients with Philadelphia chromosome positive CML in chronic phase from 2 to less than 18 years of age have been established (see sections 4.8, 5.1 and 5.2). There is no experience in paediatric patients below 2 years of age or in paediatric patients with Philadelphia chromosome positive CML in accelerated phase or blast crisis. There are no data in newly diagnosed paediatric patients below 10 years of age and limited data in imatinib-resistant or intolerant paediatric patients below 6 years of age.

Method of administration

Tasigna should be taken twice daily approximately 12 hours apart and must not be taken with food. The hard capsules should be swallowed whole with water. No food should be consumed for 2 hours before the dose is taken and no food should be consumed for at least one hour after the dose is taken.

For patients who are unable to swallow hard capsules, the content of each hard capsule may be dispersed in one teaspoon of apple sauce (puréed apple) and should be taken immediately. Not more than one teaspoon of apple sauce and no food other than apple sauce must be used (see sections 4.4 and 5.2).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Myelosuppression

Treatment with nilotinib is associated with (National Cancer Institute Common Toxicity Criteria grade 3-4) thrombocytopenia, neutropenia and anaemia. Occurrence is more frequent in patients with imatinib-resistant or intolerant CML, in particular in patients with accelerated-phase CML. Complete blood counts should be performed every two weeks for the first 2 months and then monthly thereafter, or as clinically indicated. Myelosuppression was generally reversible and usually managed by withholding Tasigna temporarily or dose reduction (see section 4.2).

QT prolongation

Nilotinib has been shown to prolong cardiac ventricular repolarisation as measured by the QT interval on the surface ECG in a concentration-dependent manner in adult and paediatric patients.

In the Phase III study in patients with newly diagnosed CML in chronic phase receiving 300 mg nilotinib twice daily, the change from baseline in mean time-averaged QTcF interval at steady state was 6 msec. No patient had a QTcF >480 msec. No episodes of torsade de pointes were observed.

In the Phase II study in imatinib-resistant and intolerant CML patients in chronic and accelerated phase receiving 400 mg nilotinib twice daily, the change from baseline in mean time-averaged QTcF interval at steady state was 5 and 8 msec, respectively. QTcF of >500 msec was observed in <1% of these patients. No episodes of torsade de pointes were observed in clinical studies.

In a healthy volunteer study with exposures that were comparable to the exposures observed in patients, the time-averaged mean placebo-subtracted QTcF change from baseline was 7 msec (CI \pm 4 msec). No subject had a QTcF >450 msec. Additionally, no clinically relevant arrhythmias were observed during the conduct of the trial. In particular, no episodes of torsade de pointes (transient or sustained) were observed.

Significant prolongation of the QT interval may occur when nilotinib is inappropriately taken with strong CYP3A4 inhibitors and/or medicinal products with a known potential to prolong QT, and/or food (see section 4.5). The presence of hypokalaemia and hypomagnesaemia may further enhance this effect. Prolongation of the QT interval may expose patients to the risk of fatal outcome.

Tasigna should be used with caution in patients who have or who are at significant risk of developing prolongation of QTc, such as those:

- with congenital long QT prolongation
- with uncontrolled or significant cardiac disease including recent myocardial infarction, congestive heart failure, unstable angina or clinically significant bradycardia.
- taking anti-arrhythmic medicinal products or other substances that lead to QT prolongation.

Close monitoring for an effect on the QTc interval is advisable and a baseline ECG is recommended prior to initiating nilotinib therapy and as clinically indicated. Hypokalaemia or hypomagnesaemia must be corrected prior to Tasigna administration and should be monitored periodically during therapy.

Sudden death

Uncommon cases (0.1 to 1%) of sudden deaths have been reported in patients with imatinib-resistant or intolerant CML in chronic phase or accelerated phase with a past medical history of cardiac disease or significant cardiac risk factors. Co-morbidities in addition to the underlying malignancy were also frequently present as were concomitant medicinal products. Ventricular repolarisation abnormalities may have been contributory factors. No cases of sudden death were reported in the Phase III study in newly diagnosed patients with CML in chronic phase.

Fluid retention and oedema

Severe forms of fluid retention such as pleural effusion, pulmonary oedema, and pericardial effusion were uncommonly (0.1 to 1%) observed in a Phase III study of newly diagnosed CML patients. Similar events were observed in post-marketing reports. Unexpected, rapid weight gain should be carefully investigated. If signs of severe fluid retention appear during treatment with nilotinib, the aetiology should be evaluated and patients treated accordingly (see section 4.2 for instructions on managing non-haematological toxicities).

Cardiovascular events

Cardiovascular events were reported in a randomised Phase III study in newly diagnosed CML patients and observed in post-marketing reports. In this clinical study with a median on-therapy time of 60.5 months, Grade 3–4 cardiovascular events included peripheral arterial occlusive disease (1.4% and 1.1% at 300 mg and 400 mg nilotinib twice daily, respectively), ischaemic heart disease (2.2% and 6.1% at 300 mg and 400 mg nilotinib twice daily, respectively) and ischaemic cerebrovascular events (1.1% and 2.2% at 300 mg and 400 mg nilotinib twice daily, respectively). Patients should be advised to seek immediate medical attention if they experience acute signs or symptoms of cardiovascular events. The cardiovascular status of patients should be evaluated and cardiovascular risk factors monitored and actively managed during nilotinib therapy according to standard guidelines. Appropriate therapy should be prescribed to manage cardiovascular risk factors (see section 4.2 for instructions on managing non-haematological toxicities).

Hepatitis B reactivation

Reactivation of hepatitis B in patients who are chronic carriers of this virus has occurred after these patients received BCR-ABL tyrosine kinase inhibitors. Some cases resulted in acute hepatic failure or fulminant hepatitis leading to liver transplantation or a fatal outcome.

Patients should be tested for HBV infection before initiating treatment with nilotinib. Experts in liver disease and in the treatment of hepatitis B should be consulted before treatment is initiated in patients with positive hepatitis B serology (including those with active disease) and for patients who test positive for HBV infection during treatment. Carriers of HBV who require treatment with nilotinib should be closely monitored for signs and symptoms of active HBV infection throughout therapy and for several months following termination of therapy (see section 4.8).

Special monitoring of Ph+ CML patients in chronic phase who have achieved a sustained deep molecular response

Eligibility for discontinuation of treatment

Eligible patients who are confirmed to express the typical BCR-ABL transcripts, e13a2/b2a2 or e14a2/b3a2, can be considered for treatment discontinuation. Patients must have typical BCR-ABL transcripts to allow quantitation of BCR-ABL, evaluation of the depth of molecular response, and determination of a possible loss of molecular remission after discontinuation of treatment with nilotinib.

Monitoring of patients who have discontinued therapy

Frequent monitoring of BCR-ABL transcript levels in patients eligible for treatment discontinuation must be performed with a quantitative diagnostic test validated to measure molecular response levels with a sensitivity of at least MR4.5 (BCR-ABL/ABL \leq 0.0032% IS). BCR-ABL transcript levels must be assessed prior to and during treatment discontinuation (see sections 4.2 and 5.1).

Loss of major molecular response (MMR=BCR-ABL/ABL \leq 0.1%IS) or confirmed loss of MR4 (two consecutive measures separated by at least 4 weeks showing loss of MR4 (MR4=BCR-ABL/ABL \leq 0.01%IS)) will trigger treatment re-initiation within 4 weeks of when loss of remission is known to have occurred. Molecular relapse can occur during the treatment-free phase, and long-term outcome data are not yet available. It is therefore crucial to perform frequent monitoring of BCR-ABL

transcript levels and complete blood count with differential in order to detect possible loss of remission (see section 4.2). For patients who fail to achieve MMR after three months of treatment re-initiation, BCR-ABL kinase domain mutation testing should be performed.

Laboratory tests and monitoring

Blood lipids

In a Phase III study in newly diagnosed CML patients, 1.1% of the patients treated with 400 mg nilotinib twice daily showed a Grade 3-4 elevation in total cholesterol; no Grade 3-4 elevations were however observed in the 300 mg twice daily dose group (see section 4.8). It is recommended that the lipid profiles be determined before initiating treatment with nilotinib, assessed at month 3 and 6 after initiating therapy and at least yearly during chronic therapy (see section 4.2). If a HMG-CoA reductase inhibitor (a lipid-lowering agent) is required, please refer to section 4.5 before initiating treatment since certain HMG-CoA reductase inhibitors are also metabolised by the CYP3A4 pathway.

Blood glucose

In a Phase III study in newly diagnosed CML patients, 6.9% and 7.2% of the patients treated with 400 mg nilotinib and 300 mg nilotinib twice daily, respectively, showed a Grade 3-4 elevation in blood glucose. It is recommended that the glucose levels be assessed before initiating treatment with Tasigna and monitored during treatment, as clinically indicated (see section 4.2). If test results warrant therapy, physicians should follow their local standards of practice and treatment guidelines.

Interactions with other medicinal products

The administration of Tasigna with agents that are strong CYP3A4 inhibitors (including, but not limited to, ketoconazole, itraconazole, voriconazole, clarithromycin, telithromycin, ritonavir) should be avoided. Should treatment with any of these agents be required, it is recommended that nilotinib therapy be interrupted if possible (see section 4.5). If transient interruption of treatment is not possible, close monitoring of the individual for prolongation of the QT interval is indicated (see sections 4.2, 4.5 and 5.2).

Concomitant use of nilotinib with medicinal products that are potent inducers of CYP3A4 (e.g. phenytoin, rifampicin, carbamazepine, phenobarbital and St. John's Wort) is likely to reduce exposure to nilotinib to a clinically relevant extent. Therefore, in patients receiving nilotinib, co-administration of alternative therapeutic agents with less potential for CYP3A4 induction should be selected (see section 4.5).

Food effect

The bioavailability of nilotinib is increased by food. Tasigna must not be taken in conjunction with food (see sections 4.2 and 4.5) and should be taken 2 hours after a meal. No food should be consumed for at least one hour after the dose is taken. Grapefruit juice and other foods that are known to inhibit CYP3A4 should be avoided. For patients who are unable to swallow hard capsules, the content of each hard capsule may be dispersed in one teaspoon of apple sauce and should be taken immediately. Not more than one teaspoon of apple sauce and no food other than apple sauce must be used (see section 5.2).

Hepatic impairment

Hepatic impairment has a modest effect on the pharmacokinetics of nilotinib. Single dose administration of 200 mg of nilotinib resulted in increases in AUC of 35%, 35% and 19% in subjects with mild, moderate and severe hepatic impairment, respectively, compared to a control group of subjects with normal hepatic function. The predicted steady-state C_{max} of nilotinib showed an increase of 29%, 18% and 22%, respectively. Clinical studies have excluded patients with alanine transaminase (ALT) and/or aspartate transaminase (AST) >2.5 (or >5, if related to disease) times the upper limit of the normal range and/or total bilirubin >1.5 times the upper limit of the normal range. Metabolism of nilotinib is mainly hepatic. Patients with hepatic impairment might therefore have increased exposure

to nilotinib and should be treated with caution (see section 4.2).

Serum lipase

Elevation in serum lipase has been observed. Caution is recommended in patients with previous history of pancreatitis. In case lipase elevations are accompanied by abdominal symptoms, nilotinib therapy should be interrupted and appropriate diagnostic measures considered to exclude pancreatitis.

Total gastrectomy

The bioavailability of nilotinib might be reduced in patients with total gastrectomy (see section 5.2). More frequent follow-up of these patients should be considered.

Tumour lysis syndrome

Due to possible occurrence of tumour lysis syndrome (TLS) correction of clinically significant dehydration and treatment of high uric acid levels are recommended prior to initiating nilotinib therapy (see section 4.8).

Lactose

Tasigna hard capsules contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Paediatric population

Laboratory abnormalities of mild to moderate transient elevations of aminotransferases and total bilirubin have been observed in children at a higher frequency than in adults, indicating a higher risk of hepatotoxicity in the paediatric population (see section 4.8). Liver function (bilirubin and hepatic transaminases levels) should be monitored monthly or as clinically indicated. Elevations of bilirubin and hepatic transaminases should be managed by withholding nilotinib temporarily, dose reduction and/or discontinuation of nilotinib (see section 4.2). The long-term effects of prolonged treatment with nilotinib in children and adolescents are unknown.

4.5 Interaction with other medicinal products and other forms of interaction

Tasigna may be given in combination with haematopoietic growth factors such as erythropoietin or granulocyte colony-stimulating factor (G-CSF) if clinically indicated. It may be given with hydroxyurea or anagrelide if clinically indicated.

Nilotinib is mainly metabolised in the liver and is also a substrate for the multi-drug efflux pump, P-glycoprotein (P-gp). Therefore, absorption and subsequent elimination of systemically absorbed nilotinib may be influenced by substances that affect CYP3A4 and/or P-gp.

Substances that may increase nilotinib serum concentrations

Concomitant administration of nilotinib with imatinib (a substrate and moderator of P-gp and CYP3A4), had a slight inhibitory effect on CYP3A4 and/or P-gp. The AUC of imatinib was increased by 18% to 39%, and the AUC of nilotinib was increased by 18% to 40%. These changes are unlikely to be clinically important.

The exposure to nilotinib in healthy subjects was increased 3-fold when co-administered with the strong CYP3A4 inhibitor ketoconazole. Concomitant treatment with strong CYP3A4 inhibitors, including ketoconazole, itraconazole, voriconazole, ritonavir, clarithromycin, and telithromycin, should therefore be avoided (see section 4.4). Increased exposure to nilotinib might also be expected with moderate CYP3A4 inhibitors. Alternative concomitant medicinal products with no or minimal CYP3A4 inhibition should be considered.

Substances that may decrease nilotinib serum concentrations

Rifampicin, a potent CYP3A4 inducer, decreases nilotinib C_{max} by 64% and reduces nilotinib AUC by 80%. Rifampicin and nilotinib should not be used concomitantly.

The concomitant administration of other medicinal products that induce CYP3A4 (e.g. phenytoin, carbamazepine, phenobarbital and St. John's Wort) is likewise likely to reduce exposure to nilotinib to a clinically relevant extent. In patients for whom CYP3A4 inducers are indicated, alternative agents with less enzyme induction potential should be selected.

Nilotinib has pH dependent solubility, with lower solubility at higher pH. In healthy subjects receiving esomeprazole at 40 mg once daily for 5 days, gastric pH was markedly increased, but nilotinib absorption was only decreased modestly (27% decrease in C_{max} and 34% decrease in AUC_{0-∞}). Nilotinib may be used concurrently with esomeprazole or other proton pump inhibitors as needed.

In a healthy subjects study, no significant change in nilotinib pharmacokinetics was observed when a single 400 mg dose of nilotinib was administered 10 hours after and 2 hours before famotidine. Therefore, when the concurrent use of a H2 blocker is necessary, it may be administered approximately 10 hours before and approximately 2 hours after the dose of Tasisna.

In the same study as above, administration of an antacid (aluminium hydroxide/magnesium hydroxide/simethicone) 2 hours before or after a single 400 mg dose of nilotinib also did not alter nilotinib pharmacokinetics. Therefore, if necessary, an antacid may be administered approximately 2 hours before or approximately 2 hours after the dose of Tasisna.

Substances that may have their systemic concentration altered by nilotinib

In vitro, nilotinib is a relatively strong inhibitor of CYP3A4, CYP2C8, CYP2C9, CYP2D6 and UGT1A1, with K_i value being lowest for CYP2C9 ($K_i=0.13$ microM).

A single-dose drug-drug interaction study in healthy volunteers with 25 mg warfarin, a sensitive CYP2C9 substrate, and 800 mg nilotinib did not result in any changes in warfarin pharmacokinetic parameters or warfarin pharmacodynamics measured as prothrombin time (PT) and international normalised ratio (INR). There are no steady-state data. This study suggests that a clinically meaningful drug-drug interaction between nilotinib and warfarin is less likely up to a dose of 25 mg of warfarin. Due to lack of steady-state data, control of warfarin pharmacodynamic markers (INR or PT) following initiation of nilotinib therapy (at least during the first 2 weeks) is recommended.

In CML patients, nilotinib administered at 400 mg twice daily for 12 days increased the systemic exposure (AUC and C_{max}) of oral midazolam (a substrate of CYP3A4) 2.6-fold and 2.0-fold, respectively. Nilotinib is a moderate CYP3A4 inhibitor. As a result, the systemic exposure of other medicinal products primarily metabolised by CYP3A4 (e.g. certain HMG-CoA reductase inhibitors) may be increased when co-administered with nilotinib. Appropriate monitoring and dose adjustment may be necessary for medicinal products that are CYP3A4 substrates and have a narrow therapeutic index (including but not limited to alfentanil, cyclosporine, dihydroergotamine, ergotamine, fentanyl, sirolimus and tacrolimus) when co-administered with nilotinib.

Anti-arrhythmic medicinal products and other substances that may prolong the QT interval

Nilotinib should be used with caution in patients who have or may develop prolongation of the QT interval, including those patients taking anti-arrhythmic medicinal products such as amiodarone, disopyramide, procainamide, quinidine and sotalol or other medicinal products that may lead to QT prolongation such as chloroquine, halofantrine, clarithromycin, haloperidol, methadone and moxifloxacin (see section 4.4).

Food interactions

The absorption and bioavailability of nilotinib are increased if it is taken with food, resulting in a higher serum concentration (see sections 4.2, 4.4 and 5.2). Grapefruit juice and other foods that are known to inhibit CYP3A4 should be avoided.

Paediatric population

Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception

Women of childbearing potential have to use highly effective contraception during treatment with nilotinib and for up to two weeks after ending treatment.

Pregnancy

There are no or limited amount of data from the use of nilotinib in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Tasigna should not be used during pregnancy unless the clinical condition of the woman requires treatment with nilotinib. If it is used during pregnancy, the patient must be informed of the potential risk to the foetus.

If a woman who is being treated with nilotinib is considering pregnancy, treatment discontinuation may be considered based on the eligibility criteria for discontinuing treatment as described in sections 4.2 and 4.4. There is a limited amount of data on pregnancies in patients while attempting treatment-free remission (TFR). If pregnancy is planned during the TFR phase, the patient must be informed of a potential need to re-initiate nilotinib treatment during pregnancy (see sections 4.2 and 4.4).

Breast-feeding

It is unknown whether nilotinib is excreted in human milk. Available toxicological data in animals have shown excretion of nilotinib in milk (see section 5.3). A risk to the newborns/infants cannot be excluded. Tasigna should not be used during breast-feeding.

Fertility

Animal studies did not show an effect on fertility in male and female rats (see section 5.3).

4.7 Effects on ability to drive and use machines

Tasigna has no or negligible influence on the ability to drive and use machines. However, it is recommended that patients experiencing dizziness, fatigue, visual impairment or other undesirable effects with a potential impact on the ability to drive or use machines safely should refrain from these activities as long as the undesirable effects persist (see section 4.8).

4.8 Undesirable effects

Summary of the safety profile

The data described below reflect exposure to nilotinib in a total of 717 adult patients from a randomised Phase III study in patients with newly diagnosed Ph+ CML in chronic phase treated at the recommended dose of 300 mg twice daily (n=279) and from an open-label multicentre Phase II study in adult patients with imatinib-resistant or intolerant CML in chronic phase (n=321) and accelerated phase (n=137) treated at the recommended dose of 400 mg twice daily. Safety information from two

Tasigna treatment discontinuation studies is also provided.

In adult patients with newly diagnosed CML in chronic phase

The median duration of exposure was 60.5 months (range 0.1-70.8 months).

The most frequent ($\geq 10\%$) non-haematological adverse reactions were rash, pruritus, headache, nausea, fatigue, alopecia, myalgia and upper abdominal pain. Most of these adverse reactions were mild to moderate in severity. Constipation, dry skin, asthenia, muscle spasms, diarrhoea, arthralgia, abdominal pain, vomiting and peripheral oedema were observed less commonly ($<10\%$ and $\geq 5\%$) were of mild to moderate severity, manageable and generally did not require dose reduction.

Treatment-emergent haematological toxicities include myelosuppression: thrombocytopenia (18%), neutropenia (15%) and anaemia (8%). Biochemical adverse drug reactions include alanine aminotransferase increased (24%), hyperbilirubinaemia (16%), aspartate aminotransferase increased (12%), lipase increased (11%), blood bilirubin increased (10%), hyperglycaemia (4%), hypercholesterolaemia (3%) and hypertriglyceridaemia ($<1\%$). Pleural and pericardial effusions, regardless of causality, occurred in 2% and $<1\%$ of patients, respectively, receiving nilotinib 300 mg twice daily. Gastrointestinal haemorrhage, regardless of causality, was reported in 3% of these patients.

The change from baseline in mean time-averaged QTcF interval at steady state was 6 msec. No patient had an absolute QTcF >500 msec while on the study medicinal product. QTcF increase from baseline exceeding 60 msec was observed in $<1\%$ of patients while on the study medicinal product. No sudden deaths or episodes of torsade de pointes (transient or sustained) were observed. No decrease from baseline in mean left ventricular ejection fraction (LVEF) was observed at any time during treatment. No patient had a LVEF of $<45\%$ during treatment nor an absolute reduction in LVEF of more than 15%.

Discontinuation due to adverse drug reactions was observed in 10% of patients.

In adult patients with imatinib-resistant or intolerant CML in chronic phase and accelerated phase

The data described below reflect exposure to nilotinib in 458 adult patients in an open-label multicentre Phase II study in patients with imatinib-resistant or intolerant CML in chronic phase (n=321) and accelerated phase (n=137) treated at the recommended dose of 400 mg twice daily.

The most frequent ($\geq 10\%$) non-haematological drug-related adverse events were rash, pruritus, nausea, fatigue, headache, vomiting, myalgia, constipation and diarrhoea. Most of these adverse events were mild to moderate in severity. Alopecia, muscle spasms, decreased appetite, arthralgia, abdominal pain, bone pain, peripheral oedema, asthenia, upper abdominal pain, dry skin, erythema and pain in extremity were observed less commonly ($<10\%$ and $\geq 5\%$) and have been of mild to moderate severity (Grade 1 or 2). Discontinuation due to adverse drug reactions was observed in 16% of chronic phase and 10% of accelerated phase patients.

Treatment-emergent haematological toxicities include myelosuppression: thrombocytopenia (31%), neutropenia (17%) and anaemia (14%). Pleural and pericardial effusions as well as complications of fluid retention occurred in $<1\%$ of patients receiving Tasigna. Cardiac failure was observed in $<1\%$ of patients. Gastrointestinal and CNS haemorrhage were reported in 1% and $<1\%$ of patients, respectively.

QTcF exceeding 500 msec was observed in $<1\%$ of patients. No episodes of torsade de pointes (transient or sustained) were observed.

Tabulated list of adverse reactions

The adverse reactions are ranked under heading of frequency using the following convention: very common ($\geq 1/10$), common ($\geq 1/100$ to $<1/10$), uncommon ($\geq 1/1,000$ to $<1/100$), rare ($\geq 1/10,000$ to $<1/1,000$), very rare ($<1/10,000$) and not known (cannot be estimated from the available data). Within

each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Most frequently reported adverse reactions in Tasigna clinical studies

Non-haematological adverse reactions (excluding laboratory abnormalities) that are reported in at least 5% of the adult patients in Tasigna clinical studies that serve as the basis for the approved indications are shown in Table 3.

Table 3 Non-haematological adverse reactions (≥5% of all patients)*

System organ class/ Adverse reaction	Newly diagnosed CML-CP 300 mg twice daily n=279			Imatinib-resistant or intolerant CML-CP and CML-AP 400 mg twice daily n=458				
	60-month analysis			24-month analysis				
	Frequency	All grades	Grade 3-4	Frequency	All grades	Grade 3-4	CML-CP n=321 Grade 3-4	CML-AP n=137 Grade 3-4
	%	%		%	%	%	%	
Metabolism and nutrition disorders								
Decreased appetite **	Common	4	0	Common	8	<1	<1	0
Nervous system disorders								
Headache	Very common	16	2	Very common	15	1	2	<1
Gastrointestinal disorders								
Nausea	Very common	14	<1	Very common	20	<1	<1	<1
Constipation	Common	10	0	Very common	12	<1	<1	0
Diarrhoea	Common	9	<1	Very common	11	2	2	<1
Vomiting	Common	6	0	Very common	10	<1	<1	0
Upper abdominal pain	Very common	10	1	Common	5	<1	<1	0
Abdominal pain	Common	6	0	Common	6	<1	<1	<1
Dyspepsia	Common	5	0	Common	3	0	0	0
Skin and subcutaneous tissue disorders								
Rash	Very common	33	<1	Very common	28	1	2	0
Pruritus	Very common	18	<1	Very common	24	<1	<1	0
Alopecia	Very common	10	0	Common	9	0	0	0
Dry skin	Common	10	0	Common	5	0	0	0
Erythema	Common	3	0	Common	5	<1	<1	0
Musculoskeletal and connective tissue disorders								
Myalgia	Very common	10	<1	Very common	10	<1	<1	<1
Muscle spasms	Common	9	0	Common	8	<1	<1	0
Arthralgia	Common	8	<1	Common	7	<1	1	0
Bone pain	Common	4	0	Common	6	<1	<1	0
Pain in	Common	5	<1	Common	5	<1	<1	<1

extremity								
General disorders and administration site conditions								
Fatigue	Very common	12	0	Very common	17	1	1	<1
Asthenia	Common	9	<1	Common	6	<1	0	<1
Oedema peripheral	Common	5	0	Common	6	0	0	0

* Percentages are rounded to integer for presentation in this table. However, percentages with one decimal precision are used to identify terms with a frequency of at least 5% and to classify terms according to frequency categories.

**Also includes preferred term anorexia

The following adverse reactions were reported in adult patients in the Tasigna clinical studies which serve as a basis for the approved indications at a frequency of less than 5%. For laboratory abnormalities, very common adverse reactions not included in Table 3 are also reported. These adverse reactions are included based on clinical relevance.

Infections and infestations

Common: folliculitis, upper respiratory tract infection (including pharyngitis, nasopharyngitis, rhinitis).

Uncommon: pneumonia, urinary tract infection, gastroenteritis, bronchitis, herpes virus infection, candidiasis (including oral candidiasis).

Not known: sepsis, subcutaneous abscess, anal abscess, furuncle, tinea pedis, hepatitis B reactivation.

Neoplasms benign, malignant and unspecified (including cysts and polyps)

Common: skin papilloma.

Not known: oral papilloma, paraproteinaemia.

Blood and lymphatic system disorders

Common: leukopenia, eosinophilia, febrile neutropenia, pancytopenia, lymphopenia.

Uncommon: thrombocythaemia, leukocytosis.

Immune system disorders

Not known: hypersensitivity.

Endocrine disorders

Uncommon: hyperthyroidism, hypothyroidism.

Not known: hyperparathyroidism secondary, thyroiditis.

Metabolism and nutrition disorders

Very common: hypophosphataemia (including blood phosphorus decreased).

Common: electrolyte imbalance (including hypomagnesaemia, hyperkalaemia, hypokalaemia, hyponatraemia, hypocalcaemia, hypercalcaemia, hyperphosphataemia), diabetes mellitus, hyperglycaemia, hypercholesterolaemia, hyperlipidaemia, hypertriglyceridaemia.

Uncommon: dehydration, increased appetite, gout, dyslipidaemia.

Not known: hyperuricaemia, hypoglycaemia.

Psychiatric disorders

Common: depression, insomnia, anxiety.

Not known: disorientation, confusional state, amnesia, dysphoria.

Nervous system disorders

Common: dizziness, peripheral neuropathy, hypoesthesia, paraesthesia.

Uncommon: intracranial haemorrhage, ischaemic stroke, transient ischaemic attack, cerebral infarction, migraine, loss of consciousness (including syncope), tremor, disturbance in attention, hyperaesthesia.

Not known: cerebrovascular accident, brain oedema, optic neuritis, lethargy, dysaesthesia, restless legs

syndrome.

Eye disorders

Common: eye haemorrhage, periorbital oedema, eye pruritus, conjunctivitis, dry eye (including xerophthalmia).

Uncommon: visual impairment, vision blurred, conjunctival haemorrhage, visual acuity reduced, eyelid oedema, photopsia, hyperaemia (scleral, conjunctival, ocular), eye irritation.

Not known: papilloedema, chorioretinopathy, diplopia, photophobia, eye swelling, blepharitis, eye pain, conjunctivitis allergic, ocular surface disease.

Ear and labyrinth disorders

Common: vertigo.

Not known: hearing impaired, ear pain, tinnitus.

Cardiac disorders

Common: angina pectoris, arrhythmia (including atroventricular block, cardiac flutter, extrasystoles, tachycardia, atrial fibrillation, bradycardia), palpitations, electrocardiogram QT prolonged.

Uncommon: cardiac failure, myocardial infarction, coronary artery disease, cardiac murmur, pericardial effusion, cyanosis.

Not known: ventricular dysfunction, pericarditis, ejection fraction decreased.

Vascular disorders

Common: hypertension, flushing, peripheral artery stenosis.

Uncommon: hypertensive crisis, peripheral arterial occlusive disease, intermittent claudication, arterial stenosis limb, haematoma, arteriosclerosis.

Not known: shock haemorrhagic, hypotension, thrombosis.

Respiratory, thoracic and mediastinal disorders

Common: dyspnoea, dyspnoea exertional, epistaxis, cough, dysphonia.

Uncommon: pulmonary oedema, pleural effusion, interstitial lung disease, pleuritic pain, pleurisy, pharyngolaryngeal pain, throat irritation.

Not known: pulmonary hypertension, wheezing, oropharyngeal pain.

Gastrointestinal disorders

Common: pancreatitis, abdominal discomfort, abdominal distension, dysgeusia, flatulence.

Uncommon: gastrointestinal haemorrhage, melaena, mouth ulceration, gastroesophageal reflux, stomatitis, oesophageal pain, dry mouth, gastritis, sensitivity of teeth.

Not known: gastrointestinal ulcer perforation, retroperitoneal haemorrhage, haematemesis, gastric ulcer, oesophagitis ulcerative, subileus, enterocolitis, haemorrhoids, hiatus hernia, rectal haemorrhage, gingivitis.

Hepatobiliary disorders

Very common: hyperbilirubinaemia (including blood bilirubin increased).

Common: hepatic function abnormal.

Uncommon: hepatotoxicity, toxic hepatitis, jaundice.

Not known: cholestasis, hepatomegaly.

Skin and subcutaneous tissue disorders

Common: night sweats, eczema, urticaria, hyperhidrosis, contusion, acne, dermatitis (including allergic, exfoliative and acneiform).

Uncommon: exfoliative rash, drug eruption, skin pain, ecchymosis, swelling face.

Not known: erythema multiforme, erythema nodosum, skin ulcer, palmar-plantar erythrodysesthesia syndrome, petechiae, photosensitivity, blister, dermal cysts, sebaceous hyperplasia, skin atrophy, skin discolouration, skin exfoliation, skin hyperpigmentation, skin hypertrophy, hyperkeratosis, psoriasis.

Musculoskeletal and connective tissue disorders

Common: musculoskeletal chest pain, musculoskeletal pain, back pain, flank pain, neck pain,

muscular weakness.

Uncommon: musculoskeletal stiffness, joint swelling.

Not known: arthritis.

Renal and urinary disorders

Common: pollakiuria.

Uncommon: dysuria, micturition urgency, nocturia.

Not known: renal failure, haematuria, urinary incontinence, chromaturia.

Reproductive system and breast disorders

Uncommon: breast pain, gynaecomastia, erectile dysfunction.

Not known: breast induration, menorrhagia, nipple swelling.

General disorders and administration site conditions

Common: chest pain (including non-cardiac chest pain), pain, pyrexia, chest discomfort, malaise.

Uncommon: face oedema, gravitational oedema, influenza-like illness, chills, feeling body temperature change (including feeling hot, feeling cold).

Not known: localised oedema.

Investigations

Very common: alanine aminotransferase increased, aspartate aminotransferase increased, lipase increased, lipoprotein cholesterol (including low density and high density) increased, total cholesterol increased, blood triglycerides increased.

Common: haemoglobin decreased, blood amylase increased, blood alkaline phosphatase increased, gamma-glutamyltransferase increased, blood creatinine phosphokinase increased, weight decreased, weight increased, blood insulin increased, globulins decreased.

Uncommon: blood lactate dehydrogenase increased, blood glucose decreased, blood urea increased.

Not known: troponin increased, blood bilirubin unconjugated increased, blood insulin decreased, insulin C-peptide decreased, blood parathyroid hormone increased.

Clinically relevant or severe abnormalities of routine haematological or biochemistry laboratory values in adult patients are presented in Table 4.

Table 4 Grade 3-4 laboratory abnormalities*

	Newly diagnosed CML-CP 300 mg twice daily	Imatinib-resistant or intolerant CML-CP and CML-AP 400 mg twice daily	
	n=279 (%)	CML-CP n=321 (%)	CML-AP n=137 (%)
Haematological parameters			
Myelosuppression			
- Neutropenia	12	31	42
- Thrombocytopenia	10	30	42
- Anaemia	4	11	27
Biochemistry parameters			
- Elevated creatinine	0	1	<1
- Elevated lipase	9	18	18
- Elevated SGOT (AST)	1	3	2
- Elevated SGPT (ALT)	4	4	4
- Hypophosphataemia	7	17	15
- Elevated bilirubin (total)	4	7	9
- Elevated glucose	7	12	6
- Elevated cholesterol (total)	0	**	**
- Elevated triglycerides	0	**	**

*Percentages with one decimal precision are used and rounded to integer for presentation in this table

**Parameters not collected

Treatment discontinuation in Ph+ CML patients in chronic phase who have achieved a sustained deep molecular response

After discontinuation of nilotinib therapy within the framework of attempting TFR, patients may experience musculoskeletal symptoms more frequently than before treatment discontinuation, e.g., myalgia, pain in extremity, arthralgia, bone pain, spinal pain or musculoskeletal pain.

In a Phase II clinical study with newly diagnosed patients with Ph+ CML in chronic phase (N=190), musculoskeletal symptoms were reported within a year of Tasigna discontinuation in 24.7% versus 16.3% within the previous year on nilotinib treatment.

In a Phase II clinical study with patients with Ph+ CML in chronic phase on nilotinib treatment and previously treated with imatinib (N=126), musculoskeletal symptoms were reported within a year of discontinuation in 42.1% versus 14.3% within the previous year on nilotinib treatment.

Description of selected adverse reactions

Sudden death

Uncommon cases (0.1 to 1%) of sudden deaths have been reported in Tasigna clinical trials and/or compassionate use programs in patients with imatinib-resistant or intolerant CML in chronic phase or accelerated phase with a past medical history of cardiac disease or significant cardiac risk factors (see section 4.4).

Hepatitis B reactivation

Hepatitis B reactivation has been reported in association with BCR-ABL TKIs. Some cases resulted in acute hepatic failure or fulminant hepatitis leading to liver transplantation or a fatal outcome (see section 4.4).

Post-marketing experience

The following adverse reactions have been derived from post-marketing experience with Tasigna via spontaneous case reports, literature cases, expanded access programmes, and clinical studies other than the global registration trials. Since these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to nilotinib exposure.

Frequency rare: Cases of tumour lysis syndrome have been reported in patients treated with nilotinib.

Paediatric population

The safety of nilotinib in paediatric patients (from 2 to <18 years of age) with Philadelphia chromosome positive CML in chronic phase (n=69) has been investigated in two studies (see section 5.1). In paediatric patients, the frequency, type and severity of adverse reactions observed have been generally consistent with those observed in adults, with the exception of the laboratory abnormalities hyperbilirubinaemia (Grade 3/4: 13.0%) and transaminase elevation (AST Grade 3/4: 1.4%, ALT Grade 3/4: 8.7%) which were reported at a higher frequency than in adult patients. Bilirubin and hepatic transaminase levels should be monitored during treatment (see sections 4.2 and 4.4).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

Isolated reports of intentional overdose with nilotinib were reported, where an unspecified number of Tasigna hard capsules were ingested in combination with alcohol and other medicinal products. Events included neutropenia, vomiting and drowsiness. No ECG changes or hepatotoxicity were reported. Outcomes were reported as recovered.

In the event of overdose, the patient should be observed and appropriate supportive treatment given.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, protein kinase inhibitors, ATC code: L01XE08

Mechanism of action

Nilotinib is a potent inhibitor of the ABL tyrosine kinase activity of the BCR-ABL oncoprotein both in cell lines and in primary Philadelphia-chromosome positive leukaemia cells. The substance binds with high affinity to the ATP-binding site in such a manner that it is a potent inhibitor of wild-type BCR-ABL and maintains activity against 32/33 imatinib-resistant mutant forms of BCR-ABL. As a consequence of this biochemical activity, nilotinib selectively inhibits the proliferation and induces apoptosis in cell lines and in primary Philadelphia-chromosome positive leukaemia cells from CML patients. In murine models of CML, as a single agent nilotinib reduces tumour burden and prolongs survival following oral administration.

Pharmacodynamic effects

Nilotinib has little or no effect against the majority of other protein kinases examined, including Src, except for the PDGF, KIT and Ephrin receptor kinases, which it inhibits at concentrations within the range achieved following oral administration at therapeutic doses recommended for the treatment of CML (see Table 5).

Table 5 Kinase profile of nilotinib (phosphorylation IC₅₀ nM)

BCR-ABL	PDGFR	KIT
20	69	210

Clinical efficacy

Clinical studies in newly diagnosed CML in chronic phase

An open-label, multicentre, randomised Phase III study was conducted to determine the efficacy of nilotinib versus imatinib in 846 adult patients with cytogenetically confirmed newly diagnosed Philadelphia chromosome positive CML in the chronic phase. Patients were within six months of diagnosis and were previously untreated, with the exception of hydroxyurea and/or anagrelide. Patients were randomised 1:1:1 to receive either nilotinib 300 mg twice daily (n=282), nilotinib 400 mg twice daily (n=281) or imatinib 400 mg once daily (n=283). Randomisation was stratified by Sokal risk score at the time of diagnosis.

Baseline characteristics were well balanced between the three treatment arms. Median age was 47 years in both nilotinib arms and 46 years in the imatinib arm, with 12.8%, 10.0% and 12.4% of patients were ≥65 years of age in the nilotinib 300 mg twice daily, nilotinib 400 mg twice daily and imatinib 400 mg once daily treatment arms, respectively. There were slightly more male than female patients (56.0%, 62.3% and 55.8%, in the nilotinib 300 mg twice daily, 400 mg twice daily and imatinib 400 mg once daily arm, respectively). More than 60% of all patients were Caucasian and 25% of all patients were Asian.

The primary data analysis time point was when all 846 patients completed 12 months of treatment (or discontinued earlier). Subsequent analyses reflect when patients completed 24, 36, 48, 60 and 72 months of treatment (or discontinued earlier). The median time on treatment was approximately 70 months in the nilotinib treatment groups and 64 months in the imatinib group. The median actual dose intensity was 593 mg/day for nilotinib 300 mg twice daily, 772 mg/day for nilotinib 400 mg twice daily and 400 mg/day for imatinib 400 mg once daily. This study is ongoing.

The primary efficacy endpoint was major molecular response (MMR) at 12 months. MMR was defined as ≤0.1% BCR-ABL/ABL% by international scale (IS) measured by RQ-PCR, which corresponds to a ≥3 log reduction of BCR-ABL transcript from standardised baseline. The MMR rate at 12 months was statistically significantly higher for nilotinib 300 mg twice daily compared to imatinib 400 mg once daily (44.3% versus 22.3%, p<0.0001). The rate of MMR at 12 months, was also statistically significantly higher for nilotinib 400 mg twice daily compared to imatinib 400 mg once daily (42.7% versus 22.3%, p<0.0001).

The rates of MMR at 3, 6, 9 and 12 months were 8.9%, 33.0%, 43.3% and 44.3% for nilotinib 300 mg twice daily, 5.0%, 29.5%, 38.1% and 42.7% for nilotinib 400 mg twice daily and 0.7%, 12.0%, 18.0% and 22.3% for imatinib 400 mg once daily.

The MMR rate at 12, 24, 36, 48, 60 and 72 months is presented in Table 6.

Table 6 MMR rate

	Nilotinib 300 mg twice daily n=282 (%)	Nilotinib 400 mg twice daily n=281 (%)	Imatinib 400 mg once daily n=283 (%)
MMR at 12 months			
Response (95% CI)	44.3 ¹ (38.4; 50.3)	42.7 ¹ (36.8; 48.7)	22.3 (17.6; 27.6)
MMR at 24 months			
Response (95% CI)	61.7 ¹ (55.8; 67.4)	59.1 ¹ (53.1; 64.9)	37.5 (31.8; 43.4)
MMR at 36 months²			
Response (95% CI)	58.5 ¹ (52.5; 64.3)	57.3 ¹ (51.3; 63.2)	38.5 (32.8; 44.5)
MMR at 48 months³			
Response (95% CI)	59.9 ¹ (54.0; 65.7)	55.2 (49.1; 61.1)	43.8 (38.0; 49.8)
MMR at 60 months⁴			
Response (95% CI)	62.8 (56.8; 68.4)	61.2 (55.2; 66.9)	49.1 (43.2; 55.1)
MMR at 72 months⁵			
Response (95% CI)	52.5 (46.5; 58.4)	57.7 (51.6; 63.5)	41.7 (35.9; 47.7)

¹ Cochran-Mantel-Haenszel (CMH) test p-value for response rate (vs. imatinib 400 mg) <0.0001

² Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 199 (35.2%) of all patients were not evaluable for MMR at 36 months (87 in the nilotinib 300 mg twice daily group and 112 in the imatinib group) due to missing/unevaluable PCR assessments (n=17), atypical transcripts at baseline (n=7), or discontinuation prior to the 36-month time point (n=175).


³ Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 305 (36.1%) of all patients were not evaluable for MMR at 48 months (98 in the nilotinib 300 mg BID group, 88 in the nilotinib 400 mg BID group and 119 in the imatinib group) due to missing/unevaluable PCR assessments (n=18), atypical transcripts at baseline (n=8), or discontinuation prior to the 48-month time point (n=279).

⁴ Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 322 (38.1%) of all patients were not evaluable for MMR at 60 months (99 in the nilotinib 300 mg twice daily group, 93 in the nilotinib 400 mg twice daily group and 130 in the imatinib group) due to missing/unevaluable PCR assessments (n=9), atypical transcripts at baseline (n=8) or discontinuation prior to the 60-month time point (n=305).

⁵ Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 395 (46.7%) of all patients were not evaluable for MMR at 72 months (130 in the nilotinib 300 mg twice daily group, 110 in the nilotinib 400 mg twice daily group and 155 in the imatinib group) due to missing/unevaluable PCR assessments (n=25), atypical transcripts at baseline (n=8) or discontinuation prior to the 72-month time point (n=362).

MMR rates by different time points (including patients who achieved MMR at or before those time points as responders) are presented in the cumulative incidence of MMR (see Figure 1).

Figure 1 Cumulative incidence of MMR



For all Sokal risk groups, the MMR rates at all time points remained consistently higher in the two nilotinib groups than in the imatinib group.

In a retrospective analysis, 91% (234/258) of patients on nilotinib 300 mg twice daily achieved BCR-ABL levels $\leq 10\%$ at 3 months of treatment compared to 67% (176/264) of patients on imatinib 400 mg once daily. Patients with BCR-ABL levels $\leq 10\%$ at 3 months of treatment show a greater overall survival at 72 months compared to those who did not achieve this molecular response level (94.5% vs. 77.1% respectively [p=0.0005]).

Based on the Kaplan-Meier analysis of time to first MMR the probability of achieving MMR at different time points was higher for both nilotinib at 300 mg and 400 mg twice daily compared to imatinib 400 mg once daily (HR=2.17 and stratified log-rank p<0.0001 between nilotinib 300 mg twice daily and imatinib 400 mg once daily, HR=1.88 and stratified log-rank p<0.0001 between nilotinib 400 mg twice daily and imatinib 400 mg once daily).

The proportion of patients who had a molecular response of $\leq 0.01\%$ and $\leq 0.0032\%$ by IS at different time points are presented in Table 7 and the proportion of patients who had a molecular response of $\leq 0.01\%$ and $\leq 0.0032\%$ by IS by different time points are presented in Figures 2 and 3. Molecular responses of $\leq 0.01\%$ and $\leq 0.0032\%$ by IS correspond to a ≥ 4 log reduction and ≥ 4.5 log reduction, respectively, of BCR-ABL transcripts from a standardised baseline.

Table 7 Proportions of patients who had molecular response of $\leq 0.01\%$ (4 log reduction) and $\leq 0.0032\%$ (4.5 log reduction)

	Nilotinib 300 mg twice daily n=282 (%)		Nilotinib 400 mg twice daily n=281 (%)		Imatinib 400 mg once daily n=283 (%)	
	$\leq 0.01\%$	$\leq 0.0032\%$	$\leq 0.01\%$	$\leq 0.0032\%$	$\leq 0.01\%$	$\leq 0.0032\%$
At 12 months	11.7	4.3	8.5	4.6	3.9	0.4
At 24 months	24.5	12.4	22.1	7.8	10.2	2.8
At 36 months	29.4	13.8	23.8	12.1	14.1	8.1
At 48 months	33.0	16.3	29.9	17.1	19.8	10.2
At 60 months	47.9	32.3	43.4	29.5	31.1	19.8
At 72 months	44.3	31.2	45.2	28.8	27.2	18.0

Figure 2 Cumulative incidence of molecular response of $\leq 0.01\%$ (4-log reduction)





Figure 3 Cumulative incidence of molecular response of $\leq 0.0032\%$ (4.5 log reduction)



Based on Kaplan-Meier estimates of the duration of first MMR, the proportions of patients who were maintaining response for 72 months among patients who achieved MMR were 92.5% (95% CI: 88.6-96.4%) in the nilotinib 300 mg twice daily group, 92.2% (95% CI: 88.5-95.9%) in the nilotinib 400 mg twice daily group and 88.0% (95% CI: 83.0-93.1%) in the imatinib 400 mg once daily group.

Complete cytogenetic response (CCyR) was defined as 0% Ph+ metaphases in the bone marrow based on a minimum of 20 metaphases evaluated. Best CCyR rate by 12 months (including patients who achieved CCyR at or before the 12 month time point as responders) was statistically higher for both nilotinib 300 mg and 400 mg twice daily compared to imatinib 400 mg once daily, see Table 8.

CCyR rate by 24 months (includes patients who achieved CCyR at or before the 24 month time point as responders) was statistically higher for both the nilotinib 300 mg twice daily and 400 mg twice daily groups compared to the imatinib 400 mg once daily group.

Table 8 Best CCyR rate

	Nilotinib 300 mg twice daily n=282 (%)	Nilotinib 400 mg twice daily n=281 (%)	Imatinib 400 mg once daily n=283 (%)
By 12 months			
Response (95% CI)	80.1 (75.0; 84.6)	77.9 (72.6; 82.6)	65.0 (59.2; 70.6)
No response	19.9	22.1	35.0
CMH test p-value for response rate (versus imatinib 400 mg once daily)	<0.0001	0.0005	
By 24 months			
Response (95% CI)	86.9 (82.4; 90.6)	84.7 (79.9; 88.7)	77.0 (71.7; 81.8)
No response	13.1	15.3	23.0
CMH test p-value for response rate (versus imatinib 400 mg once daily)	0.0018	0.0160	

Based on Kaplan-Meier estimates, the proportions of patients who were maintaining response for 72 months among patients who achieved CCyR were 99.1% (95% CI: 97.9-100%) in the nilotinib 300 mg twice daily group, 98.7% (95% CI: 97.1-100%) in the nilotinib 400 mg twice daily group and 97.0% (95% CI: 94.7-99.4%) in the imatinib 400 mg once daily group.

Progression to accelerated phase (AP) or blast crisis (BC) on treatment is defined as the time from the date of randomisation to the first documented disease progression to accelerated phase or blast crisis or CML-related death. Progression to accelerated phase or blast crisis on treatment was observed in a total of 17 patients: 2 patients on nilotinib 300 mg twice daily, 3 patients on nilotinib 400 mg twice daily and 12 patients on imatinib 400 mg once daily. The estimated rates of patients free from progression to accelerated phase or blast crisis at 72 months were 99.3%, 98.7% and 95.2%, respectively (HR=0.1599 and stratified log-rank p=0.0059 between nilotinib 300 mg twice daily and imatinib once daily, HR=0.2457 and stratified log-rank p=0.0185 between nilotinib 400 mg twice daily and imatinib once daily). No new events of progression to AP/BC were reported on-treatment since the 2-year analysis.

Including clonal evolution as a criterion for progression, a total of 25 patients progressed to accelerated phase or blast crisis on treatment by the cut-off date (3 in the nilotinib 300 mg twice daily group, 5 in the nilotinib 400 mg twice daily group and 17 in the imatinib 400 mg once daily group). The estimated rates of patients free from progression to accelerated phase or blast crisis including clonal evolution at 72 months were 98.7%, 97.9% and 93.2%, respectively (HR=0.1626 and stratified log-rank p=0.0009 between nilotinib 300 mg twice daily and imatinib once daily, HR=0.2848 and stratified log-rank p=0.0085 between nilotinib 400 mg twice daily and imatinib once daily).

A total of 55 patients died during treatment or during the follow-up after discontinuation of treatment. (21 in the nilotinib 300 mg twice daily group, 11 in the nilotinib 400 mg twice daily group and 23 in the imatinib 400 mg once daily group). Twenty-six (26) of these 55 deaths were related to CML (6 in the nilotinib 300 mg twice daily group, 4 in the nilotinib 400 mg twice daily group and 16 in the imatinib 400 mg once daily group). The estimated rates of patients alive at 72 months were 91.6%, 95.8% and 91.4%, respectively (HR=0.8934 and stratified log-rank p=0.7085 between nilotinib 300 mg twice daily and imatinib, HR=0.4632 and stratified log-rank p=0.0314 between nilotinib 400 mg twice daily and imatinib). Considering only CML-related deaths as events, the estimated rates of overall survival at 72 months were 97.7%, 98.5% and 93.9%, respectively (HR=0.3694 and stratified log-rank p=0.0302 between nilotinib 300 mg twice daily and imatinib, HR=0.2433 and stratified log-rank p=0.0061 between nilotinib 400 mg twice daily and imatinib).

Clinical studies in imatinib-resistant or intolerant CML in chronic phase and accelerated phase

An open-label, uncontrolled, multicentre Phase II study was conducted to determine the efficacy of nilotinib in adult patients with imatinib resistant or intolerant CML with separate treatment arms for chronic and accelerated phase disease. Efficacy was based on 321 CP patients and 137 AP patients enrolled. Median duration of treatment was 561 days for CP patients and 264 days for AP patients (see Table 9). Tasigna was administered on a continuous basis (twice daily 2 hours after a meal and with no food for at least one hour after administration) unless there was evidence of inadequate response or disease progression. The dose was 400 mg twice daily and dose escalation to 600 mg twice daily was allowed.

Table 9 Duration of exposure with nilotinib

	Chronic phase n=321	Accelerated phase n=137
Median duration of therapy in days (25th-75th percentiles)	561 (196-852)	264 (115-595)

Resistance to imatinib included failure to achieve a complete haematological response (by 3 months), cytogenetic response (by 6 months) or major cytogenetic response (by 12 months) or progression of disease after a previous cytogenetic or haematological response. Imatinib intolerance included patients who discontinued imatinib because of toxicity and were not in major cytogenetic response at time of study entry.

Overall, 73% of patients were imatinib-resistant, while 27% were imatinib-intolerant. The majority of patients had a long history of CML that included extensive prior treatment with other antineoplastic agents, including imatinib, hydroxyurea, interferon, and some had even failed organ transplant (Table 10). The median highest prior imatinib dose had been 600 mg/day. The highest prior imatinib dose was ≥ 600 mg/day in 74% of all patients, with 40% of patients receiving imatinib doses ≥ 800 mg/day.

Table 10 CML disease history characteristics

	Chronic phase (n=321)	Accelerated phase (n=137)*
Median time since diagnosis in months (range)	58 (5-275)	71 (2-298)
Imatinib Resistant	226 (70%)	109 (80%)
Intolerant without MCyR	95 (30%)	27 (20%)
Median time of imatinib treatment in days (25th-75 th percentiles)	975 (519-1,488)	857 (424-1,497)
Prior hydroxyurea	83%	91%
Prior interferon	58%	50%
Prior bone marrow transplant	7%	8%

* Missing information on imatinib-resistant/intolerant status for one patient.

The primary endpoint in the CP patients was major cytogenetic response (MCyR), defined as elimination (CCyR, complete cytogenetic response) or significant reduction to $<35\%$ Ph+ metaphases (partial cytogenetic response) of Ph+ haematopoietic cells. Complete haematological response (CHR) in CP patients was evaluated as a secondary endpoint. The primary endpoint in the AP patients was overall confirmed haematological response (HR), defined as either a complete haematological response, no evidence of leukaemia or return to chronic phase.

Chronic phase

The MCyR rate in 321 CP patients was 51%. Most responders achieved their MCyR rapidly within

3 months (median 2.8 months) of starting nilotinib treatment and these were sustained. The median time to achieve CCyR was just past 3 months (median 3.4 months). Of the patients who achieved MCyR, 77% (95% CI: 70% - 84%) were maintaining response at 24 months. Median duration of MCyR has not been reached. Of the patients who achieved CCyR, 85% (95% CI: 78% - 93%) were maintaining response at 24 months. Median duration of CCyR has not been reached. Patients with a CHR at baseline achieved a MCyR faster (1.9 versus 2.8 months). Of CP patients without a baseline CHR, 70% achieved a CHR, median time to CHR was 1 month and median duration of CHR was 32.8 months. The estimated 24-month overall survival rate in CML-CP patients was 87%.

Accelerated phase

The overall confirmed HR rate in 137 AP patients was 50%. Most responders achieved a HR early with nilotinib treatment (median 1.0 months) and these have been durable (median duration of confirmed HR was 24.2 months). Of the patients who achieved HR, 53% (95% CI: 39% - 67%) were maintaining response at 24 months. MCyR rate was 30% with a median time to response of 2.8 months. Of the patients who achieved MCyR, 63% (95% CI: 45% - 80%) were maintaining response at 24 months. Median duration of MCyR was 32.7 months. The estimated 24-month overall survival rate in CML-AP patients was 70%.

The rates of response for the two treatment arms are reported in Table 11.

Table 11 Response in CML

(Best response rate)	Chronic phase			Accelerated phase		
	Intolerant (n=95)	Resistant (n=226)	Total (n=321)	Intolerant (n=27)	Resistant (n=109)	Total* (n=137)
Haematological Response (%)						
Overall (95% CI)	-	-	-	48	51	50
Complete	87	65	70 ¹	(29-68)	(42-61)	(42-59)
NEL	(74-94)	(56-72)	(63-76)	37	28	30
Return to CP	-	-	-	7	10	9
	-	-		4	13	11
Cytogenetic Response (%)						
Major (95% CI)	57	49	51 (46-57)	33	29	30
Complete	(46-67)	(42-56)	37	(17-54)	(21-39)	(22-38)
Partial	41	35	15	22	19	20
	16	14		11	10	10

NEL = no evidence of leukaemia/marrow response

¹ 114 CP patients had a CHR at baseline and were therefore not assessable for complete haematological response

* Missing information on imatinib-resistant/intolerant status for one patient.

Efficacy data in patients with CML-BC are not yet available. Separate treatment arms were also included in the Phase II study to investigate Tasigna in a group of CP and AP patients who had been extensively pre-treated with multiple therapies including a tyrosine kinase inhibitor agent in addition to imatinib. Of these patients 30/36 (83%) were treatment resistant not intolerant. In 22 CP patients evaluated for efficacy nilotinib induced a 32% MCyR rate and a 50% CHR rate. In 11 AP patients, evaluated for efficacy, treatment induced a 36% overall HR rate.

After imatinib failure, 24 different BCR-ABL mutations were noted in 42% of chronic phase and 54% of accelerated phase CML patients who were evaluated for mutations. Tasigna demonstrated efficacy in patients harboring a variety of BCR-ABL mutations associated with imatinib resistance, except T315I.

Treatment discontinuation in Ph+ CML patients in chronic phase who have been treated with nilotinib as first-line therapy and who have achieved a sustained deep molecular response

In an open-label, single-arm study, 215 adult patients with Ph+ CML in chronic phase treated with nilotinib in first-line for ≥ 2 years who achieved MR4.5 as measured with the MolecularMD MRDx BCR-ABL test were enrolled to continue nilotinib treatment for additional 52 weeks (nilotinib consolidation phase). 190 of 215 patients (88.4%) entered the TFR phase after achieving a sustained deep molecular response during the consolidation phase, defined by the following criteria:

- the 4 last quarterly assessments (taken every 12 weeks) were at least MR4 (BCR-ABL/ABL $\leq 0.01\%$ IS), and maintained for one year
- the last assessment being MR4.5 (BCR-ABL/ABL $\leq 0.0032\%$ IS)
- no more than two assessments falling between MR4 and MR4.5 ($0.0032\% \text{ IS} < \text{BCR-ABL/ABL} \leq 0.01\% \text{ IS}$).


The primary endpoint was the percentage of patients in MMR at 48 weeks after starting the TFR phase (considering any patient who required re-initiation of treatment as non-responder). Of the 190 patients who entered the TFR phase, 98 patients (51.6% [95% CI: 44.2, 58.9]) were in MMR at 48 weeks.

Eighty-eight patients (46.3%) discontinued the TFR phase due to loss of MMR, and 1 (0.5%), 1 (0.5%), and 3 patients (1.6%) due to death from unknown cause, physician decision and subject decision, respectively. Among these 88 patients, 86 patients restarted nilotinib treatment and 2 patients permanently discontinued the study. Eighty-five of these 86 patients (98.8%) regained MMR, (one patient discontinued study permanently due to subject decision) and 76 patients (88.4%) regained MR4.5 by the time of the cut-off date.

The Kaplan-Meier (KM) estimated median time on nilotinib treatment to regain MMR and MR4.5 was 7.9 weeks (95% CI: 5.1, 8.0) and 13.1 weeks (95% CI: 12.3, 15.7), respectively. The KM estimated MMR and MR4.5 rates at 24 weeks of re-initiation were 98.8 % (95% CI: 94.2, 99.9) and 90.9 % (95% CI: 83.2, 96.0), respectively.

The KM estimate of median treatment-free survival (TFS) has not yet been reached (Figure 4); 99 of 190 patients (52.1%) did not have a TFS event.

Figure 4 Kaplan-Meier estimate of treatment-free survival after start of TFR (full analysis set)



Treatment discontinuation in CML patients in chronic phase who have achieved a sustained deep

molecular response on nilotinib treatment following prior imatinib therapy

In an open-label, single-arm study, 163 adult patients with Ph+ CML in chronic phase taking tyrosine kinase inhibitors (TKIs) for ≥ 3 years (imatinib as initial TKI therapy for more than 4 weeks without documented MR4.5 on imatinib at the time of switch to nilotinib, then switched to nilotinib for at least two years), and who achieved MR4.5 on nilotinib treatment as measured with the MolecularMD MRDx BCR-ABL test were enrolled to continue nilotinib treatment for additional 52 weeks (nilotinib consolidation phase). 126 of 163 patients (77.3%) entered the TFR phase after achieving a sustained deep molecular response during the consolidation phase, defined by the following criterion:

- The 4 last quarterly assessments (taken every 12 weeks) showed no confirmed loss of MR4.5 (BCR-ABL/ABL $\leq 0.0032\%$ IS) during one year.


The primary endpoint was the proportion of patients without confirmed loss of MR4.0 or loss of MMR within 48 weeks following treatment discontinuation. Of the 126 patients who entered the TFR phase, 73 patients (57.9%, [95% CI: 48.8, 66.7]) had no loss of MMR, no confirmed loss of MR4.0, and no re-initiation of nilotinib within 48 weeks.

Among the 53 patients who discontinued the TFR phase due to confirmed loss of MR4.0 or loss of MMR, 51 patients restarted nilotinib and 2 patients discontinued the study. Forty-eight of these 51 patients (94.1%) regained MR4.0 and 47 patients (92.2%) regained MR4.5 by the time of the cut-off date.

The Kaplan-Meier (KM) estimated median time on nilotinib to regain MR4.0 and MR4.5 was 12.0 weeks (95% CI: 8.3, 12.7) and 13.1 weeks (95% CI: 12.4, 16.1), respectively. The KM estimated MR4.0 and MR4.5 rates at 48 weeks of re-initiation were 100.0% (95% CI: not estimated) and 94.8% (95% CI: 85.1, 99.0), respectively.

The median TFS has not yet been reached (Figure 5); 74 of 126 patients (58.7%) did not have a TFS event.

Figure 5 Kaplan-Meier estimate of treatment-free survival after start of TFR (full analysis set)



Paediatric population

The safety and efficacy of nilotinib in paediatric patients with Ph+ CML in chronic phase have been investigated in two studies. A total of 69 paediatric patients (from 2 to <18 years of age) with either newly diagnosed Ph+ CML in chronic phase (n=25) or imatinib/dasatinib resistant or imatinib-intolerant Ph+ CML in chronic phase (n=44) received nilotinib treatment at a dose of 230 mg/m² twice

daily, rounded to the nearest 50 mg dose (to a maximum single dose of 400 mg).

In the pooled CML patient population, the median actual dose intensity was 435.5 mg/m²/day (range: 149 to 517 mg/m²/day), and the median relative dose intensity was 94.7% (range: 32 to 112%). Forty patients (58.0%) had relative dose intensity superior to 90%. The median time on treatment with nilotinib was 13.80 months (range: 0.7-30.9 months).

In the resistant or intolerant CML patients, the major molecular response (MMR; BCR-ABL/ABL ≤0.1% IS) rate was 40.9% (95% CI: 26.3, 56.8) at 12 cycles, with 18 patients being in MMR. In the newly diagnosed CML patients, the MMR rate was 60.0% (95% CI: 38.7, 78.9) at 12 cycles, with 15 patients achieving MMR. In resistant or intolerant CML patients, the cumulative MMR rate was 47.7% by cycle 12. In newly diagnosed CML patients, the cumulative MMR rate was 64.0% by cycle 12.

Among the 21 resistant or intolerant CML patients who were in MMR at any time on treatment, the median time to first MMR was 2.76 months (95% CI: 0.03, 5.55). For the 17 newly diagnosed CML patients who achieved MMR, the median time to first MMR was 5.55 months (95% CI: 5.52, 5.75).

Among resistant or intolerant CML patients, the percentage of patients who achieved BCR-ABL/ABL ≤0.01% IS (MR4.0) by the cut-off date was 11.4%, while 4.5% of the patients achieved BCR-ABL/ABL ≤0.0032% IS (MR4.5). Among newly diagnosed patients, the percentage of patients who achieved MR4.0 was 32%, while 28.0% achieved MR4.5.

None of the 21 resistant or intolerant CML patients who were in MMR on treatment had confirmed loss of MMR. Among the 17 newly diagnosed CML patients who achieved MMR, one patient had confirmed loss of MMR (the patient lost CHR due to an increase in basophil count, however did not progress to AP/BC).

One resistant or intolerant CML patient progressed to AP/BC after about 10 months on treatment.

No deaths were reported on treatment or after treatment discontinuation in both studies.

5.2 Pharmacokinetic properties

Absorption

Peak concentrations of nilotinib are reached 3 hours after oral administration. Nilotinib absorption following oral administration was approximately 30%. The absolute bioavailability of nilotinib has not been determined. As compared to an oral drink solution (pH of 1.2 to 1.3), relative bioavailability of nilotinib capsule is approximately 50%. In healthy volunteers, C_{max} and area under the serum concentration-time curve (AUC) of nilotinib are increased by 112% and 82%, respectively, compared to fasting conditions when Tasigna is given with food. Administration of Tasigna 30 minutes or 2 hours after food increased bioavailability of nilotinib by 29% or 15%, respectively (see sections 4.2, 4.4 and 4.5).

Nilotinib absorption (relative bioavailability) might be reduced by approximately 48% and 22% in patients with total gastrectomy and partial gastrectomy, respectively.

Distribution

The blood-to-plasma ratio of nilotinib is 0.71. Plasma protein binding is approximately 98% on the basis of *in vitro* experiments.

Biotransformation

Main metabolic pathways identified in healthy subjects are oxidation and hydroxylation. Nilotinib is the main circulating component in the serum. None of the metabolites contribute significantly to the

pharmacological activity of nilotinib. Nilotinib is primarily metabolised by CYP3A4, with possible minor contribution from CYP2C8.

Elimination

After a single dose of radiolabelled nilotinib in healthy subjects, more than 90% of the dose was eliminated within 7 days, mainly in faeces (94% of the dose). Unchanged nilotinib accounted for 69% of the dose.

The apparent elimination half-life estimated from the multiple-dose pharmacokinetics with daily dosing was approximately 17 hours. Inter-patient variability in nilotinib pharmacokinetics was moderate to high.

Linearity/non-linearity

Steady-state nilotinib exposure was dose-dependent, with less than dose-proportional increases in systemic exposure at dose levels higher than 400 mg given as once-daily dosing. Daily systemic exposure to nilotinib with 400 mg twice-daily dosing at steady state was 35% higher than with 800 mg once-daily dosing. Systemic exposure (AUC) of nilotinib at steady state at a dose level of 400 mg twice daily was approximately 13.4% higher than at a dose level of 300 mg twice daily. The average nilotinib trough and peak concentrations over 12 months were approximately 15.7% and 14.8% higher following 400 mg twice-daily dosing compared to 300 mg twice daily. There was no relevant increase in exposure to nilotinib when the dose was increased from 400 mg twice daily to 600 mg twice daily.

Steady-state conditions were essentially achieved by day 8. An increase in serum exposure to nilotinib between the first dose and steady state was approximately 2-fold for daily dosing and 3.8-fold for twice-daily dosing.

Bioavailability/bioequivalence studies

Single-dose administration of 400 mg nilotinib, using 2 hard capsules of 200 mg whereby the content of each hard capsule was dispersed in one teaspoon of apple sauce, was shown to be bioequivalent with a single-dose administration of 2 intact hard capsules of 200 mg.

Paediatric population

Following administration of nilotinib in paediatric patients at 230 mg/m² twice daily, rounded to the nearest 50 mg dose (to a maximum single dose of 400 mg), steady-state exposure and clearance of nilotinib were found to be similar (within 2-fold) to adult patients treated with 400 mg twice daily. The pharmacokinetic exposure of nilotinib following a single or multiple doses appeared to be comparable between paediatric patients from 2 years to <10 years and from ≥10 years to <18 years.

5.3 Preclinical safety data

Nilotinib has been evaluated in safety pharmacology, repeated dose toxicity, genotoxicity, reproductive toxicity, phototoxicity and carcinogenicity (rats and mice) studies.

Nilotinib did not have effects on CNS or respiratory functions. *In vitro* cardiac safety studies demonstrated a preclinical signal for QT prolongation, based upon block of hERG currents and prolongation of the action potential duration in isolated rabbit hearts by nilotinib. No effects were seen in ECG measurements in dogs or monkeys treated for up to 39 weeks or in a special telemetry study in dogs.

Repeated-dose toxicity studies in dogs of up to 4 weeks' duration and in cynomolgus monkeys of up to 9 months' duration revealed the liver as the primary target organ of toxicity of nilotinib. Alterations included increased alanine aminotransferase and alkaline phosphatase activity and histopathology findings (mainly sinusoidal cell or Kupffer cell hyperplasia/hypertrophy, bile duct hyperplasia and

periportal fibrosis). In general the changes in clinical chemistry were fully reversible after a four-week recovery period and the histological alterations showed partial reversibility. Exposures at the lowest dose levels at which the liver effects were seen were lower than the exposure in humans at a dose of 800 mg/day. Only minor liver alterations were seen in mice or rats treated for up to 26 weeks. Mainly reversible increases in cholesterol levels were seen in rats, dogs and monkeys.

Genotoxicity studies in bacterial *in vitro* systems and in mammalian *in vitro* and *in vivo* systems with and without metabolic activation did not reveal any evidence for a mutagenic potential of nilotinib.

In the 2-year rat carcinogenicity study, the major target organ for non-neoplastic lesions was the uterus (dilatation, vascular ectasia, endothelial cell hyperplasia, inflammation and/or epithelial hyperplasia). There was no evidence of carcinogenicity upon administration of nilotinib at 5, 15 and 40 mg/kg/day. Exposures (in terms of AUC) at the highest dose level represented approximately 2x to 3x human daily steady-state exposure (based on AUC) to nilotinib at the dose of 800 mg/day.

In the 26-week Tg.rasH2 mouse carcinogenicity study, in which nilotinib was administered at 30, 100 and 300 mg/kg/day, skin papillomas/carcinomas were detected at 300 mg/kg, representing approximately 30 to 40 times (based on AUC) the human exposure at the maximum approved dose of 800 mg/day (administered as 400 mg twice daily). The No-Observed-Effect-Level for the skin neoplastic lesions was 100 mg/kg/day, representing approximately 10 to 20 times the human exposure at the maximum approved dose of 800 mg/day (administered as 400 mg twice daily). The major target organs for non-neoplastic lesions were the skin (epidermal hyperplasia), the growing teeth (degeneration/atrophy of the enamel organ of upper incisors and inflammation of the gingiva/odontogenic epithelium of incisors) and the thymus (increased incidence and/or severity of decreased lymphocytes).

Nilotinib did not induce teratogenicity, but did show embryo- and foetotoxicity at doses that also showed maternal toxicity. Increased post-implantation loss was observed in both the fertility study, which involved treatment of both males and females, and the embryotoxicity study, which involved treatment of females. Embryo-lethality and foetal effects (mainly decreased foetal weights, premature fusion of the facial bones (fused maxilla/zygomatic) visceral and skeletal variations) in rats and increased resorption of foetuses and skeletal variations in rabbits were present in the embryotoxicity studies. In a pre- and postnatal development study in rats, maternal exposure to nilotinib caused reduced pup body weight with associated changes in physical development parameters as well as reduced mating and fertility indices in the offspring. Exposure to nilotinib in females at No-Observed-Adverse-Effect-Levels was generally less or equal to that in humans at 800 mg/day.

In a juvenile development study, nilotinib was administered via oral gavage to juvenile rats from the first week post partum through young adult (day 70 post partum) at doses of 2, 6 and 20 mg/kg/day. Besides standard study parameters, evaluations of developmental landmarks, CNS effects, mating and fertility were performed. Based on a reduction in body weight in both genders and a delayed preputial separation in males (which may be associated with the reduction in weight), the No-Observed-Effect-Level in juvenile rats was considered to be 6 mg/kg/day. The juvenile animals did not exert increased sensitivity to nilotinib relative to adults. In addition, the toxicity profile in juvenile rats was comparable to that observed in adult rats.

No effects on sperm count/motility or on fertility were noted in male and female rats up to the highest tested dose, approximately 5 times the recommended dosage for humans.

Nilotinib was shown to absorb light in the UV-B and UV-A range, is distributed into the skin and showed a phototoxic potential *in vitro*, but no effects have been observed *in vivo*. Therefore the risk that nilotinib causes photosensitisation in patients is considered very low.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tasigna 50 mg hard capsules

Capsule content

Lactose monohydrate
Crospovidone Type A
Poloxamer 188
Colloidal anhydrous silica
Magnesium stearate

Capsule shell

Gelatin
Titanium dioxide (E171)
Red iron oxide (E172)
Yellow iron oxide (E172)

Printing ink

Shellac
Black iron oxide (E172)
Propylene glycol
Ammonium hydroxide

Tasigna 200 mg hard capsules

Capsule content

Lactose monohydrate
Crospovidone Type A
Poloxamer 188
Colloidal anhydrous silica
Magnesium stearate

Capsule shell

Gelatin
Titanium dioxide (E171)
Yellow iron oxide (E172)

Printing ink

Shellac (E904)
Red iron oxide (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Do not store above 30°C.

Store in the original package in order to protect from moisture.

6.5 Nature and contents of container

Tasigna is available in the following pack sizes:

Tasigna 50 mg hard capsules

PVC/PVDC/Alu blisters

- Pack containing 120 (3 packs of 40) hard capsules.

Tasigna 200 mg hard capsules

PVC/PVDC/Alu blisters

- Unit packs containing 28 hard capsules in a wallet.
- Unit packs containing 28 hard capsules (7 daily blisters, each containing 4 hard capsules) or 40 hard capsules (5 blisters, each containing 8 hard capsules).
- Multipacks containing 112 (4 wallets of 28) hard capsules.
- Multipacks containing 112 (4 packs of 28) hard capsules, 120 (3 packs of 40) hard capsules or 392 (14 packs of 28) hard capsules.

PA/Alu/PVC/Alu blisters

- Unit packs containing 28 hard capsules in a wallet.
- Multipacks containing 112 (4 packs of 28) hard capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

Tasigna 50 mg hard capsules

EU/1/07/422/015

Tasigna 200 mg hard capsules

EU/1/07/422/001-004
EU/1/07/422/007-008
EU/1/07/422/011-012
EU/1/07/422/014

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 19 November 2007
Date of latest renewal: 19 November 2012

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 150 mg hard capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One hard capsule contains 150 mg nilotinib (as hydrochloride monohydrate).

Excipient with known effect

One hard capsule contains 117.08 mg lactose monohydrate.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule.

White to yellowish powder in red opaque hard gelatin capsules, size 1 with black axial imprint "NVR/BCR".

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Tasigna is indicated for the treatment of:

- adult and paediatric patients with newly diagnosed Philadelphia chromosome positive chronic myelogenous leukaemia (CML) in the chronic phase,
- paediatric patients with Philadelphia chromosome positive CML in chronic phase with resistance or intolerance to prior therapy including imatinib.

4.2 Posology and method of administration

Therapy should be initiated by a physician experienced in the diagnosis and the treatment of patients with CML.

Posology

Treatment should be continued as long as clinical benefit is observed or until unacceptable toxicity occurs.

If a dose is missed the patient should not take an additional dose, but take the usual prescribed next dose.

Posology for Philadelphia chromosome positive CML adult patients

The recommended dose is 300 mg twice daily.

For a dose of 400 mg once daily (see dose adjustments below), 200 mg hard capsules are available.

Posology for Philadelphia chromosome positive CML paediatric patients

Dosing in paediatric patients is individualised and is based on body surface area (mg/m²). The recommended dose of nilotinib is 230 mg/m² twice daily, rounded to the nearest 50 mg dose (to a maximum single dose of 400 mg) (see Table 1). Different strengths of Tasigna hard capsules can be combined to attain the desired dose.

There is no experience with treatment of paediatric patients below 2 years of age. There are no data in newly diagnosed paediatric patients below 10 years of age and limited data in imatinib-resistant or intolerant paediatric patients below 6 years of age.

Table 1 Dosing scheme of nilotinib 230 mg/m² twice daily

Body Surface Area (BSA)	Dose in mg (twice daily)
Up to 0.32 m ²	50 mg
0.33 – 0.54 m ²	100 mg
0.55 – 0.76 m ²	150 mg
0.77 – 0.97 m ²	200 mg
0.98 – 1.19 m ²	250 mg
1.20 – 1.41 m ²	300 mg
1.42 – 1.63 m ²	350 mg
≥1.64 m ²	400 mg

Philadelphia chromosome positive CML patients in chronic phase who have been treated with nilotinib as first-line therapy and who achieved a sustained deep molecular response (MR4.5)

Discontinuation of treatment may be considered in eligible Philadelphia chromosome positive (Ph+) CML patients in chronic phase who have been treated with nilotinib at 300 mg twice daily for a minimum of 3 years if a deep molecular response is sustained for a minimum of one year immediately prior to discontinuation of therapy. Discontinuation of nilotinib therapy should be initiated by a physician experienced in the treatment of patients with CML (see sections 4.4 and 5.1).

Eligible patients who discontinue nilotinib therapy must have their BCR-ABL transcript levels and complete blood count with differential monitored monthly for one year, then every 6 weeks for the second year, and every 12 weeks thereafter. Monitoring of BCR-ABL transcript levels must be performed with a quantitative diagnostic test validated to measure molecular response levels on the International Scale (IS) with a sensitivity of at least MR4.5 (BCR-ABL/ABL ≤0.0032% IS).

For patients who lose MR4 (MR4=BCR-ABL/ABL ≤0.01%IS) but not MMR (MMR=BCR-ABL/ABL ≤0.1%IS) during the treatment-free phase, BCR-ABL transcript levels should be monitored every 2 weeks until BCR-ABL levels return to a range between MR4 and MR4.5. Patients who maintain BCR-ABL levels between MMR and MR4 for a minimum of 4 consecutive measurements can return to the original monitoring schedule.

Patients who lose MMR must re-initiate treatment within 4 weeks of when loss of remission is known to have occurred. Nilotinib therapy should be re-initiated at 300 mg twice daily or at a reduced dose level of 400 mg once daily if the patient had a dose reduction prior to discontinuation of therapy. Patients who re-initiate nilotinib therapy should have their BCR-ABL transcript levels monitored monthly until MMR is re-established and every 12 weeks thereafter (see section 4.4).

Dose adjustments or modifications

Tasigna may need to be temporarily withheld and/or dose reduced for haematological toxicities (neutropenia, thrombocytopenia) that are not related to the underlying leukaemia (see Table 2).

Table 2 Dose adjustments for neutropenia and thrombocytopenia

Adult patients with newly diagnosed chronic phase CML at 300 mg twice daily	ANC* <1.0 x 10 ⁹ /l and/or platelet counts <50 x 10 ⁹ /l	<ol style="list-style-type: none"> 1. Treatment with nilotinib must be interrupted and blood count monitored. 2. Treatment must be resumed within 2 weeks at prior dose if ANC >1.0 x 10⁹/l and/or platelets >50 x 10⁹/l. 3. If blood counts remain low, a dose reduction to 400 mg once daily may be required.
Paediatric patients with newly diagnosed chronic phase CML at 230 mg/m ² twice daily and imatinib-resistant or intolerant CML in chronic phase at 230 mg/m ² twice daily	ANC* <1.0 x 10 ⁹ /l and/or platelet counts <50 x 10 ⁹ /l	<ol style="list-style-type: none"> 1. Treatment with nilotinib must be interrupted and blood count monitored. 2. Treatment must be resumed within 2 weeks at prior dose if ANC >1.5 x 10⁹/l and/or platelets >75 x 10⁹/l. 3. If blood counts remain low, a dose reduction to 230 mg/m² once daily may be required. 4. If event occurs after dose reduction, consider discontinuing treatment.

*ANC = absolute neutrophil count

If clinically significant moderate or severe non-haematological toxicity develops, dosing should be interrupted, and patients should be monitored and treated accordingly. If the prior dose was 300 mg twice daily in adult patients or 230 mg/m² twice daily in paediatric patients, dosing may be resumed at 400 mg once daily in adult patients and at 230 mg/m² once daily in paediatric patients once the toxicity has resolved. If the prior dose was 400 mg once daily in adult patients or 230 mg/m² once daily in paediatric patients, treatment should be discontinued. If clinically appropriate, re-escalation of the dose to 300 mg twice daily in adult patients or to 230 mg/m² twice daily in paediatric patients should be considered.

Elevated serum lipase: For Grade 3-4 serum lipase elevations, doses in adult patients should be reduced to 400 mg once daily or interrupted. In paediatric patients, treatment must be interrupted until the event returns to Grade ≤1. Thereafter, if the prior dose was 230 mg/m² twice daily, treatment can be resumed at 230 mg/m² once daily. If the prior dose was 230 mg/m² once daily, treatment should be discontinued. Serum lipase levels should be tested monthly or as clinically indicated (see section 4.4).

Elevated bilirubin and hepatic transaminases: For Grade 3-4 bilirubin and hepatic transaminase elevations in adult patients, doses should be reduced to 400 mg once daily or interrupted. For Grade ≥2 bilirubin elevations or Grade ≥3 hepatic transaminase elevations in paediatric patients, treatment must be interrupted until the levels return to Grade ≤1. Thereafter, if the prior dose was 230 mg/m² twice daily, treatment can be resumed at 230 mg/m² once daily. If the prior dose was 230 mg/m² once daily, and recovery to Grade ≤1 takes longer than 28 days, treatment should be discontinued. Bilirubin and hepatic transaminases levels should be tested monthly or as clinically indicated.

Special populations

Elderly

Approximately 12% of subjects in the clinical study were 65 years of age or over. No major differences were observed for safety and efficacy in patients ≥65 years of age as compared to adults aged 18 to 65 years.

Renal impairment

Clinical studies have not been performed in patients with impaired renal function. Since nilotinib and its metabolites are not renally excreted, a decrease in total body clearance is not anticipated in patients with renal impairment.

Hepatic impairment

Hepatic impairment has a modest effect on the pharmacokinetics of nilotinib. Dose adjustment is not considered necessary in patients with hepatic impairment. However, patients with hepatic impairment should be treated with caution (see section 4.4).

Cardiac disorders

In clinical studies, patients with uncontrolled or significant cardiac disease (e.g. recent myocardial infarction, congestive heart failure, unstable angina or clinically significant bradycardia) were excluded. Caution should be exercised in patients with relevant cardiac disorders (see section 4.4).

Increases in total serum cholesterol levels have been reported with nilotinib therapy (see section 4.4). Lipid profiles should be determined prior to initiating nilotinib therapy, assessed at month 3 and 6 after initiating therapy and at least yearly during chronic therapy.

Increases in blood glucose levels have been reported with nilotinib therapy (see section 4.4). Blood glucose levels should be assessed prior to initiating nilotinib therapy and monitored during treatment.

Paediatric population

The safety and efficacy of Tasigna in paediatric patients with Philadelphia chromosome positive CML in chronic phase from 2 to less than 18 years of age have been established (see sections 4.8, 5.1 and 5.2). There is no experience in paediatric patients below 2 years of age or in paediatric patients with Philadelphia chromosome positive CML in accelerated phase or blast crisis. There are no data in newly diagnosed paediatric patients below 10 years of age and limited data in imatinib-resistant or intolerant paediatric patients below 6 years of age.

Method of administration

Tasigna should be taken twice daily approximately 12 hours apart and must not be taken with food. The hard capsules should be swallowed whole with water. No food should be consumed for 2 hours before the dose is taken and for at least one hour after.

For patients who are unable to swallow hard capsules, the content of each hard capsule may be dispersed in one teaspoon of apple sauce (puréed apple) and should be taken immediately. Not more than one teaspoon of apple sauce and no food other than apple sauce must be used (see sections 4.4 and 5.2).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Myelosuppression

Treatment with nilotinib is associated with (National Cancer Institute Common Toxicity Criteria grade 3-4) thrombocytopenia, neutropenia and anaemia. Complete blood counts should be performed every two weeks for the first 2 months and then monthly thereafter, or as clinically indicated. Myelosuppression was generally reversible and usually managed by withholding Tasigna temporarily or dose reduction (see section 4.2).

QT prolongation

Nilotinib has been shown to prolong cardiac ventricular repolarisation as measured by the QT interval on the surface ECG in a concentration-dependent manner in adult and paediatric patients.

In the Phase III study in patients with newly diagnosed CML in chronic phase receiving 300 mg nilotinib twice daily, the change from baseline in mean time-averaged QTcF interval at steady state was 6 msec. No patient had a QTcF >480 msec. No episodes of torsade de pointes were observed.

In a healthy volunteer study with exposures that were comparable to the exposures observed in patients, the time-averaged mean placebo-subtracted QTcF change from baseline was 7 msec (CI ± 4 msec). No subject had a QTcF >450 msec. Additionally, no clinically relevant arrhythmias were observed during the conduct of the trial. In particular, no episodes of torsade de pointes (transient or sustained) were observed.

Significant prolongation of the QT interval may occur when nilotinib is inappropriately taken with strong CYP3A4 inhibitors and/or medicinal products with a known potential to prolong QT, and/or food (see section 4.5). The presence of hypokalaemia and hypomagnesaemia may further enhance this effect. Prolongation of the QT interval may expose patients to the risk of fatal outcome.

Tasigna should be used with caution in patients who have or who are at significant risk of developing prolongation of QTc, such as those:

- with congenital long QT prolongation
- with uncontrolled or significant cardiac disease including recent myocardial infarction, congestive heart failure, unstable angina or clinically significant bradycardia.
- taking anti-arrhythmic medicinal products or other substances that lead to QT prolongation.

Close monitoring for an effect on the QTc interval is advisable and a baseline ECG is recommended prior to initiating nilotinib therapy and as clinically indicated. Hypokalaemia or hypomagnesaemia must be corrected prior to Tasigna administration and should be monitored periodically during therapy.

Sudden death

Uncommon cases (0.1 to 1%) of sudden deaths have been reported in patients with imatinib-resistant or intolerant CML in chronic phase or accelerated phase with a past medical history of cardiac disease or significant cardiac risk factors. Co-morbidities in addition to the underlying malignancy were also frequently present as were concomitant medicinal products. Ventricular repolarisation abnormalities may have been contributory factors. No cases of sudden death were reported in the Phase III study in newly diagnosed patients with CML in chronic phase.

Fluid retention and oedema

Severe forms of fluid retention such as pleural effusion, pulmonary oedema, and pericardial effusion were uncommonly (0.1 to 1%) observed in a Phase III study of newly diagnosed CML patients. Similar events were observed in post-marketing reports. Unexpected, rapid weight gain should be carefully investigated. If signs of severe fluid retention appear during treatment with nilotinib, the aetiology should be evaluated and patients treated accordingly (see section 4.2 for instructions on managing non-haematological toxicities).

Cardiovascular events

Cardiovascular events were reported in a randomised Phase III study in newly diagnosed CML patients and observed in post-marketing reports. In this clinical study with a median on-therapy time of 60.5 months, Grade 3-4 cardiovascular events included peripheral arterial occlusive disease (1.4% and 1.1% at 300 mg and 400 mg nilotinib twice daily, respectively), ischaemic heart disease (2.2% and 6.1% at 300 mg and 400 mg nilotinib twice daily, respectively) and ischaemic cerebrovascular events (1.1% and 2.2% at 300 mg and 400 mg nilotinib twice daily, respectively). Patients should be advised

to seek immediate medical attention if they experience acute signs or symptoms of cardiovascular events. The cardiovascular status of patients should be evaluated and cardiovascular risk factors monitored and actively managed during nilotinib therapy according to standard guidelines. Appropriate therapy should be prescribed to manage cardiovascular risk factors (see section 4.2 for instructions on managing non-haematological toxicities).

Hepatitis B reactivation

Reactivation of hepatitis B in patients who are chronic carriers of this virus has occurred after these patients received BCR-ABL tyrosine kinase inhibitors. Some cases resulted in acute hepatic failure or fulminant hepatitis leading to liver transplantation or a fatal outcome.

Patients should be tested for HBV infection before initiating treatment with nilotinib. Experts in liver disease and in the treatment of hepatitis B should be consulted before treatment is initiated in patients with positive hepatitis B serology (including those with active disease) and for patients who test positive for HBV infection during treatment. Carriers of HBV who require treatment with nilotinib should be closely monitored for signs and symptoms of active HBV infection throughout therapy and for several months following termination of therapy (see section 4.8).

Special monitoring of Ph+ CML patients in chronic phase who have achieved a sustained deep molecular response

Eligibility for discontinuation of treatment

Eligible patients who are confirmed to express the typical BCR-ABL transcripts, e13a2/b2a2 or e14a2/b3a2, can be considered for treatment discontinuation. Patients must have typical BCR-ABL transcripts to allow quantitation of BCR-ABL, evaluation of the depth of molecular response, and determination of a possible loss of molecular remission after discontinuation of treatment with nilotinib.

Monitoring of patients who have discontinued therapy

Frequent monitoring of BCR-ABL transcript levels in patients eligible for treatment discontinuation must be performed with a quantitative diagnostic test validated to measure molecular response levels with a sensitivity of at least MR4.5 ($\text{BCR-ABL/ABL} \leq 0.0032\% \text{ IS}$). BCR-ABL transcript levels must be assessed prior to and during treatment discontinuation (see sections 4.2 and 5.1).

Loss of major molecular response ($\text{MMR} = \text{BCR-ABL/ABL} \leq 0.1\% \text{ IS}$) or confirmed loss of MR4 (two consecutive measures separated by at least 4 weeks showing loss of MR4 ($\text{MR4} = \text{BCR-ABL/ABL} \leq 0.01\% \text{ IS}$)) will trigger treatment re-initiation within 4 weeks of when loss of remission is known to have occurred. Molecular relapse can occur during the treatment-free phase, and long-term outcome data are not yet available. It is therefore crucial to perform frequent monitoring of BCR-ABL transcript levels and complete blood count with differential in order to detect possible loss of remission (see section 4.2). For patients who fail to achieve MMR after three months of treatment re-initiation, BCR-ABL kinase domain mutation testing should be performed.

Laboratory tests and monitoring

Blood lipids

In a Phase III study in newly diagnosed CML patients, 1.1% of the patients treated with 400 mg nilotinib twice daily showed a Grade 3-4 elevation in total cholesterol; no Grade 3-4 elevations were however observed in the 300 mg twice daily dose group (see section 4.8). It is recommended that the lipid profiles be determined before initiating treatment with nilotinib, assessed at month 3 and 6 after initiating therapy and at least yearly during chronic therapy (see section 4.2). If a HMG-CoA reductase inhibitor (a lipid-lowering agent) is required, please refer to section 4.5 before initiating treatment since certain HMG-CoA reductase inhibitors are also metabolised by the CYP3A4 pathway.

Blood glucose

In a Phase III study in newly diagnosed CML patients, 6.9% and 7.2% of the patients treated with

400 mg nilotinib and 300 mg nilotinib twice daily, respectively, showed a Grade 3-4 elevation in blood glucose. It is recommended that the glucose levels be assessed before initiating treatment with Tasigna and monitored during treatment, as clinically indicated (see section 4.2). If test results warrant therapy, physicians should follow their local standards of practice and treatment guidelines.

Interactions with other medicinal products

The administration of Tasigna with agents that are strong CYP3A4 inhibitors (including, but not limited to, ketoconazole, itraconazole, voriconazole, clarithromycin, telithromycin, ritonavir) should be avoided. Should treatment with any of these agents be required, it is recommended that nilotinib therapy be interrupted if possible (see section 4.5). If transient interruption of treatment is not possible, close monitoring of the individual for prolongation of the QT interval is indicated (see sections 4.2, 4.5 and 5.2).

Concomitant use of nilotinib with medicinal products that are potent inducers of CYP3A4 (e.g. phenytoin, rifampicin, carbamazepine, phenobarbital and St. John's Wort) is likely to reduce exposure to nilotinib to a clinically relevant extent. Therefore, in patients receiving nilotinib, co-administration of alternative therapeutic agents with less potential for CYP3A4 induction should be selected (see section 4.5).

Food effect

The bioavailability of nilotinib is increased by food. Tasigna must not be taken in conjunction with food (see sections 4.2 and 4.5) and should be taken 2 hours after a meal. No food should be consumed for at least one hour after the dose is taken. Grapefruit juice and other foods that are known to inhibit CYP3A4 should be avoided. For patients who are unable to swallow hard capsules, the content of each hard capsule may be dispersed in one teaspoon of apple sauce and should be taken immediately. Not more than one teaspoon of apple sauce and no food other than apple sauce must be used (see section 5.2).

Hepatic impairment

Hepatic impairment has a modest effect on the pharmacokinetics of nilotinib. Single dose administration of 200 mg of nilotinib resulted in increases in AUC of 35%, 35% and 19% in subjects with mild, moderate and severe hepatic impairment, respectively, compared to a control group of subjects with normal hepatic function. The predicted steady-state C_{max} of nilotinib showed an increase of 29%, 18% and 22%, respectively. Clinical studies have excluded patients with alanine transaminase (ALT) and/or aspartate transaminase (AST) >2.5 (or >5, if related to disease) times the upper limit of the normal range and/or total bilirubin >1.5 times the upper limit of the normal range. Metabolism of nilotinib is mainly hepatic. Patients with hepatic impairment might therefore have increased exposure to nilotinib and should be treated with caution (see section 4.2).

Serum lipase

Elevation in serum lipase has been observed. Caution is recommended in patients with previous history of pancreatitis. In case lipase elevations are accompanied by abdominal symptoms, nilotinib therapy should be interrupted and appropriate diagnostic measures considered to exclude pancreatitis.

Total gastrectomy

The bioavailability of nilotinib might be reduced in patients with total gastrectomy (see section 5.2). More frequent follow-up of these patients should be considered.

Tumour lysis syndrome

Due to possible occurrence of tumour lysis syndrome (TLS) correction of clinically significant dehydration and treatment of high uric acid levels are recommended prior to initiating nilotinib

therapy (see section 4.8).

Lactose

Tasigna hard capsules contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Paediatric population

Laboratory abnormalities of mild to moderate transient elevations of aminotransferases and total bilirubin have been observed in children at a higher frequency than in adults, indicating a higher risk of hepatotoxicity in the paediatric population (see section 4.8). Liver function (bilirubin and hepatic transaminases levels) should be monitored monthly or as clinically indicated. Elevations of bilirubin and hepatic transaminases should be managed by withholding nilotinib temporarily, dose reduction and/or discontinuation of nilotinib (see section 4.2). The long-term effects of prolonged treatment with nilotinib in children and adolescents are unknown.

4.5 Interaction with other medicinal products and other forms of interaction

Tasigna may be given in combination with haematopoietic growth factors such as erythropoietin or granulocyte colony-stimulating factor (G-CSF) if clinically indicated. It may be given with hydroxyurea or anagrelide if clinically indicated.

Nilotinib is mainly metabolised in the liver and is also a substrate for the multi-drug efflux pump, P-glycoprotein (P-gp). Therefore, absorption and subsequent elimination of systemically absorbed nilotinib may be influenced by substances that affect CYP3A4 and/or P-gp.

Substances that may increase nilotinib serum concentrations

Concomitant administration of nilotinib with imatinib (a substrate and moderator of P-gp and CYP3A4), had a slight inhibitory effect on CYP3A4 and/or P-gp. The AUC of imatinib was increased by 18% to 39%, and the AUC of nilotinib was increased by 18% to 40%. These changes are unlikely to be clinically important.

The exposure to nilotinib in healthy subjects was increased 3-fold when co-administered with the strong CYP3A4 inhibitor ketoconazole. Concomitant treatment with strong CYP3A4 inhibitors, including ketoconazole, itraconazole, voriconazole, ritonavir, clarithromycin, and telithromycin, should therefore be avoided (see section 4.4). Increased exposure to nilotinib might also be expected with moderate CYP3A4 inhibitors. Alternative concomitant medicinal products with no or minimal CYP3A4 inhibition should be considered.

Substances that may decrease nilotinib serum concentrations

Rifampicin, a potent CYP3A4 inducer, decreases nilotinib C_{max} by 64% and reduces nilotinib AUC by 80%. Rifampicin and nilotinib should not be used concomitantly.

The concomitant administration of other medicinal products that induce CYP3A4 (e.g. phenytoin, carbamazepine, phenobarbital and St. John's Wort) is likewise likely to reduce exposure to nilotinib to a clinically relevant extent. In patients for whom CYP3A4 inducers are indicated, alternative agents with less enzyme induction potential should be selected.

Nilotinib has pH dependent solubility, with lower solubility at higher pH. In healthy subjects receiving esomeprazole at 40 mg once daily for 5 days, gastric pH was markedly increased, but nilotinib absorption was only decreased modestly (27% decrease in C_{max} and 34% decrease in $AUC_{0-\infty}$). Nilotinib may be used concurrently with esomeprazole or other proton pump inhibitors as needed.

In a healthy subjects study, no significant change in nilotinib pharmacokinetics was observed when a

single 400 mg dose of nilotinib was administered 10 hours after and 2 hours before famotidine. Therefore, when the concurrent use of a H2 blocker is necessary, it may be administered approximately 10 hours before and approximately 2 hours after the dose of Tasigna.

In the same study as above, administration of an antacid (aluminium hydroxide/magnesium hydroxide/simethicone) 2 hours before or after a single 400 mg dose of nilotinib also did not alter nilotinib pharmacokinetics. Therefore, if necessary, an antacid may be administered approximately 2 hours before or approximately 2 hours after the dose of Tasigna.

Substances that may have their systemic concentration altered by nilotinib

In vitro, nilotinib is a relatively strong inhibitor of CYP3A4, CYP2C8, CYP2C9, CYP2D6 and UGT1A1, with K_i value being lowest for CYP2C9 ($K_i=0.13$ microM).

A single-dose drug-drug interaction study in healthy volunteers with 25 mg warfarin, a sensitive CYP2C9 substrate, and 800 mg nilotinib did not result in any changes in warfarin pharmacokinetic parameters or warfarin pharmacodynamics measured as prothrombin time (PT) and international normalised ratio (INR). There are no steady-state data. This study suggests that a clinically meaningful drug-drug interaction between nilotinib and warfarin is less likely up to a dose of 25 mg of warfarin. Due to lack of steady-state data, control of warfarin pharmacodynamic markers (INR or PT) following initiation of nilotinib therapy (at least during the first 2 weeks) is recommended.

In CML patients, nilotinib administered at 400 mg twice daily for 12 days increased the systemic exposure (AUC and C_{max}) of oral midazolam (a substrate of CYP3A4) 2.6-fold and 2.0-fold, respectively. Nilotinib is a moderate CYP3A4 inhibitor. As a result, the systemic exposure of other medicinal products primarily metabolised by CYP3A4 (e.g. certain HMG-CoA reductase inhibitors) may be increased when co-administered with nilotinib. Appropriate monitoring and dose adjustment may be necessary for medicinal products that are CYP3A4 substrates and have a narrow therapeutic index (including but not limited to alfentanil, cyclosporine, dihydroergotamine, ergotamine, fentanyl, sirolimus and tacrolimus) when co-administered with nilotinib.

Anti-arrhythmic medicinal products and other substances that may prolong the QT interval

Nilotinib should be used with caution in patients who have or may develop prolongation of the QT interval, including those patients taking anti-arrhythmic medicinal products such as amiodarone, disopyramide, procainamide, quinidine and sotalol or other medicinal products that may lead to QT prolongation such as chloroquine, halofantrine, clarithromycin, haloperidol, methadone and moxifloxacin (see section 4.4).

Food interactions

The absorption and bioavailability of nilotinib are increased if it is taken with food, resulting in a higher serum concentration (see sections 4.2, 4.4 and 5.2). Grapefruit juice and other foods that are known to inhibit CYP3A4 should be avoided.

Paediatric population

Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception

Women of childbearing potential have to use highly effective contraception during treatment with nilotinib and for up to two weeks after ending treatment.

Pregnancy

There are no or limited amount of data from the use of nilotinib in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Tasigna should not be used during pregnancy unless the clinical condition of the woman requires treatment with nilotinib. If it is used during pregnancy, the patient must be informed of the potential risk to the foetus.

If a woman who is being treated with nilotinib is considering pregnancy, treatment discontinuation may be considered based on the eligibility criteria for discontinuing treatment as described in sections 4.2 and 4.4. There is a limited amount of data on pregnancies in patients while attempting treatment-free remission (TFR). If pregnancy is planned during the TFR phase, the patient must be informed of a potential need to re-initiate nilotinib treatment during pregnancy (see sections 4.2 and 4.4).

Breast-feeding

It is unknown whether nilotinib is excreted in human milk. Available toxicological data in animals have shown excretion of nilotinib in milk (see section 5.3). A risk to the newborns/infants cannot be excluded. Tasigna should not be used during breast-feeding.

Fertility

Animal studies did not show an effect on fertility in male and female rats (see section 5.3).

4.7 Effects on ability to drive and use machines

Tasigna has no or negligible influence on the ability to drive and use machines. However, it is recommended that patients experiencing dizziness, fatigue, visual impairment or other undesirable effects with a potential impact on the ability to drive or use machines safely should refrain from these activities as long as the undesirable effects persist (see section 4.8).

4.8 Undesirable effects

Summary of the safety profile

The data described below reflect exposure to nilotinib in 279 adult patients from a randomised Phase III study in patients with newly diagnosed Ph+ CML in chronic phase treated with 300 mg of nilotinib twice daily. Safety information from a Tasigna treatment discontinuation study in CML patients who have been treated with nilotinib as first-line therapy is also provided.

The median duration of exposure was 60.5 months (range 0.1-70.8 months).

The most frequent ($\geq 10\%$) non-haematological adverse reactions were rash, pruritus, headache, nausea, fatigue, alopecia, myalgia and upper abdominal pain. Most of these adverse reactions were mild to moderate in severity. Constipation, dry skin, asthenia, muscle spasms, diarrhoea, arthralgia, abdominal pain, vomiting and peripheral oedema were observed less commonly ($< 10\%$ and $\geq 5\%$), were of mild to moderate severity, manageable and generally did not require dose reduction.

Treatment-emergent haematological toxicities include myelosuppression: thrombocytopenia (18%), neutropenia (15%) and anaemia (8%). Biochemical adverse drug reactions include alanine aminotransferase increased (24%), hyperbilirubinaemia (16%), aspartate aminotransferase increased (12%), lipase increased (11%), blood bilirubin increased (10%), hyperglycaemia (4%), hypercholesterolaemia (3%) and hypertriglyceridaemia ($< 1\%$). Pleural and pericardial effusions, regardless of causality, occurred in 2% and $< 1\%$ of patients, respectively, receiving nilotinib 300 mg twice daily. Gastrointestinal haemorrhage, regardless of causality, was reported in 3% of these patients.

The change from baseline in mean time-averaged QTcF interval at steady state was 6 msec. No patient had an absolute QTcF >500 msec while on the study medicinal product. QTcF increase from baseline exceeding 60 msec was observed in <1% of patients while on the study medicinal product. No sudden deaths or episodes of torsade de pointes (transient or sustained) were observed. No decrease from baseline in mean left ventricular ejection fraction (LVEF) was observed at any time during treatment. No patient had a LVEF of <45% during treatment nor an absolute reduction in LVEF of more than 15%.

Discontinuation due to adverse drug reactions was observed in 10% of patients.

Tabulated list of adverse reactions

The adverse reactions are ranked under heading of frequency using the following convention: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Most frequently reported adverse reactions in Tasigna clinical studies

Non-haematological adverse reactions (excluding laboratory abnormalities) that are reported in at least 5% of the adult patients treated with 300 mg of nilotinib twice daily in the randomised Phase III study are shown in Table 3.

Table 3 Non-haematological adverse reactions ($\geq 5\%$ of all patients)*

System organ class	Frequency	Adverse reaction	All grades %	Grade 3-4 %
Nervous system disorders	Very common	Headache	16	2
Gastrointestinal disorders	Very common	Nausea	14	<1
	Very common	Abdominal pain upper	10	1
	Common	Constipation	10	0
	Common	Diarrhoea	9	<1
	Common	Abdominal pain	6	0
	Common	Vomiting	6	0
	Common	Dyspepsia	5	0
Skin and subcutaneous tissue disorders	Very common	Rash	33	<1
	Very common	Pruritus	18	<1
	Very common	Alopecia	10	0
	Common	Dry skin	10	0
Musculoskeletal and connective tissue disorders	Very common	Myalgia	10	<1
	Common	Muscle spasms	9	0
	Common	Arthralgia	8	<1
	Common	Pain in extremity	5	<1
General disorders and administration site conditions	Very common	Fatigue	12	0
	Common	Asthenia	9	<1
	Common	Oedema peripheral	5	<1

*Percentages are rounded to integer for presentation in this table. However, percentages with one decimal precision are used to identify terms with a frequency of at least 5% and to classify terms according to frequency categories.

The following adverse reactions were reported in adult patients in the Tasigna Phase III study at a frequency of less than 5%. For laboratory abnormalities, very common events ($\geq 1/10$) not included in

Table 3 are also reported. These adverse reactions are included based on clinical relevance and ranked in order of decreasing seriousness within each category using the following convention: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), not known (cannot be estimated from the available data).

Infections and infestations

Common: folliculitis, upper respiratory tract infection (including pharyngitis, nasopharyngitis, rhinitis).

Not known: herpes virus infection, oral candidiasis, subcutaneous abscess, anal abscess, tinea pedis, hepatitis B reactivation.

Neoplasms benign, malignant and unspecified (including cysts and polyps)

Common: skin papilloma.

Not known: oral papilloma, paraproteinaemia.

Blood and lymphatic system disorders

Common: leukopenia, eosinophilia, lymphopenia.

Uncommon: pancytopenia.

Not known: febrile neutropenia.

Immune system disorders

Not known: hypersensitivity.

Endocrine disorders

Not known: hyperparathyroidism secondary.

Metabolism and nutrition disorders

Very common: hypophosphataemia (including blood phosphorus decreased).

Common: diabetes mellitus, hypercholesterolaemia, hyperlipidaemia, hypertriglyceridaemia, hyperglycaemia, decreased appetite, hypocalcaemia, hypokalaemia.

Uncommon: hyperkalaemia, dyslipidaemia, gout.

Not known: hyperuricaemia, hypoglycaemia, appetite disorder.

Psychiatric disorders

Common: insomnia, depression, anxiety.

Not known: amnesia, dysphoria.

Nervous system disorders

Common: dizziness, hypoaesthesia, peripheral neuropathy.

Uncommon: ischaemic stroke, cerebral infarction, migraine, paraesthesia.

Not known: cerebrovascular accident, basilar artery stenosis, syncope, tremor, lethargy, dysaesthesia, restless legs syndrome, hyperaesthesia.

Eye disorders

Common: eye pruritus, conjunctivitis, dry eye (including xerophthalmia).

Uncommon: eyelid oedema, photopsia, conjunctival haemorrhage, hyperaemia (scleral, conjunctival, ocular).

Not known: periorbital oedema, blepharitis, eye pain, chorioretinopathy, conjunctivitis allergic, ocular surface disease, vision blurred.

Ear and labyrinth disorders

Common: vertigo.

*Cardiac disorders**

Common: angina pectoris, arrhythmia (including atrioventricular block, tachycardia, atrial fibrillation, ventricular extrasystoles, bradycardia), electrocardiogram QT prolonged, palpitations, myocardial

infarction.

Uncommon: cardiac failure, cyanosis.

Not known: ejection fraction decrease, pericardial effusion, pericarditis, diastolic dysfunction, left bundle branch block.

**reported in 300 mg twice daily and/or 400 mg twice daily treatment arm of phase III study*

Vascular disorders

Common: hypertension, flushing.

Uncommon: intermittent claudication, peripheral arterial occlusive disease, arteriosclerosis.

Not known: haematoma, peripheral artery stenosis.

Respiratory, thoracic and mediastinal disorders

Common: dyspnoea, cough.

Uncommon: pleural effusion.

Not known: dyspnoea exertional, pleurisy, epistaxis, oropharyngeal pain.

Gastrointestinal disorders

Common: abdominal distension, abdominal discomfort, dysgeusia, flatulence.

Uncommon: pancreatitis, gastritis, sensitivity of teeth.

Not known: oesophageal ulcer, gastric ulcer, oesophageal pain, stomatitis, dry mouth, enterocolitis, haemorrhoids, hiatus hernia, rectal haemorrhage, gingivitis.

Hepatobiliary disorders

Very common: hyperbilirubinaemia (including blood bilirubin increased).

Common: hepatic function abnormal.

Uncommon: jaundice.

Not known: toxic hepatitis.

Skin and subcutaneous tissue disorders

Common: erythema, hyperhidrosis, contusion, acne, dermatitis (including allergic, exfoliative and acneiform), night sweats, eczema.

Uncommon: drug eruption, skin pain.

Not known: erythema multiforme, urticaria, blister, dermal cyst, sebaceous hyperplasia, swelling face, skin atrophy, skin hypertrophy, skin exfoliation, skin hyperpigmentation, skin discolouration, hyperkeratosis, psoriasis.

Musculoskeletal and connective tissue disorders

Common: bone pain, back pain, muscular weakness.

Uncommon: musculoskeletal pain, flank pain.

Renal and urinary disorders

Not known: dysuria, pollakiuria, chromaturia.

Reproductive system and breast disorders

Uncommon: erectile dysfunction.

Not known: gynaecomastia, breast induration, menorrhagia, nipple swelling.

General disorders and administration site conditions

Common: pyrexia, chest pain (including non-cardiac chest pain), chest discomfort.

Uncommon: pain, chills, feeling body temperature change (including feeling hot, feeling cold), malaise.

Not known: face oedema, localised oedema.

Investigations

Very common: alanine aminotransferase increased, aspartate aminotransferase increased, lipase increased, lipoprotein cholesterol (including low density and high density) increased, total cholesterol

increased, blood triglycerides increased.

Common: haemoglobin decreased, blood amylase increased, blood alkaline phosphatase increased, gamma-glutamyltransferase increased, weight increased, blood insulin increased, globulins decreased.

Not known: blood parathyroid hormone increased, blood insulin decreased, insulin C-peptide decreased, weight decreased.

Clinically relevant or severe abnormalities of routine haematological or biochemistry laboratory values in adult patients are presented in Table 4.

Table 4 Grade 3-4 laboratory abnormalities*

	n=279 (%)
Haematological parameters	
Myelosuppression	
- Neutropenia	12
- Thrombocytopenia	10
- Anaemia	4
Biochemistry parameters	
- Elevated creatinine	0
- Elevated lipase	9
- Elevated SGOT (AST)	1
- Elevated SGPT (ALT)	4
- Hypophosphataemia	7
- Elevated bilirubin (total)	4
- Elevated glucose	7
- Elevated cholesterol (total)	0
- Elevated triglycerides	0

*Percentages with one decimal precision are used and rounded to integer for presentation in this table.

Treatment discontinuation in Ph+ CML patients in chronic phase who have achieved a sustained deep molecular response

After discontinuation of nilotinib therapy within the framework of attempting TFR, patients may experience musculoskeletal symptoms more frequently than before treatment discontinuation, e.g., myalgia, pain in extremity, arthralgia, bone pain, spinal pain or musculoskeletal pain.

In a Phase II clinical study with newly diagnosed patients with Ph+ CML in chronic phase (N=190), musculoskeletal symptoms were reported within a year of Tasigna discontinuation in 24.7% versus 16.3% within the previous year on nilotinib treatment.

Description of selected adverse reactions

Hepatitis B reactivation

Hepatitis B reactivation has been reported in association with BCR-ABL TKIs. Some cases resulted in acute hepatic failure or fulminant hepatitis leading to liver transplantation or a fatal outcome (see section 4.4).

Post-marketing experience

The following adverse reactions have been derived from post-marketing experience with Tasigna via spontaneous case reports, literature cases, expanded access programmes, and clinical studies other than the global registration trials. Since these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal

relationship to nilotinib exposure.

Frequency rare: Cases of tumour lysis syndrome have been reported in patients treated with nilotinib.

Paediatric population

The safety of nilotinib in paediatric patients (from 2 to <18 years of age) with Philadelphia chromosome positive CML in chronic phase (n=69) has been investigated in two studies (see section 5.1). In paediatric patients, the frequency, type and severity of adverse reactions observed have been generally consistent with those observed in adults, with the exception of the laboratory abnormalities hyperbilirubinaemia (Grade 3/4: 13.0%) and transaminase elevation (AST Grade 3/4: 1.4%, ALT Grade 3/4: 8.7%) which were reported at a higher frequency than in adult patients. Bilirubin and hepatic transaminase levels should be monitored during treatment (see sections 4.2 and 4.4).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

Isolated reports of intentional overdose with nilotinib were reported, where an unspecified number of Tasigna hard capsules were ingested in combination with alcohol and other medicinal products. Events included neutropenia, vomiting and drowsiness. No ECG changes or hepatotoxicity were reported. Outcomes were reported as recovered.

In the event of overdose, the patient should be observed and appropriate supportive treatment given.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, protein kinase inhibitors, ATC code: L01XE08

Mechanism of action

Nilotinib is a potent inhibitor of the ABL tyrosine kinase activity of the BCR-ABL oncoprotein both in cell lines and in primary Philadelphia-chromosome positive leukaemia cells. The substance binds with high affinity to the ATP-binding site in such a manner that it is a potent inhibitor of wild-type BCR-ABL and maintains activity against 32/33 imatinib-resistant mutant forms of BCR-ABL. As a consequence of this biochemical activity, nilotinib selectively inhibits the proliferation and induces apoptosis in cell lines and in primary Philadelphia-chromosome positive leukaemia cells from CML patients. In murine models of CML, as a single agent nilotinib reduces tumour burden and prolongs survival following oral administration.

Pharmacodynamic effects

Nilotinib has little or no effect against the majority of other protein kinases examined, including Src, except for the PDGF, KIT and Ephrin receptor kinases, which it inhibits at concentrations within the range achieved following oral administration at therapeutic doses recommended for the treatment of CML (see Table 5).

Table 5 Kinase profile of nilotinib (phosphorylation IC₅₀ nM)

BCR-ABL	PDGFR	KIT
20	69	210

Clinical efficacy

Clinical studies in newly diagnosed CML in chronic phase

An open-label, multicentre, randomised Phase III study was conducted to determine the efficacy of nilotinib versus imatinib in 846 adult patients with cytogenetically confirmed newly diagnosed Philadelphia chromosome positive CML in the chronic phase. Patients were within six months of diagnosis and were previously untreated, with the exception of hydroxyurea and/or anagrelide. Patients were randomised 1:1:1 to receive either nilotinib 300 mg twice daily (n=282), nilotinib 400 mg twice daily (n=281) or imatinib 400 mg once daily (n=283). Randomisation was stratified by Sokal risk score at the time of diagnosis.

Baseline characteristics were well balanced between the three treatment arms. Median age was 47 years in both nilotinib arms and 46 years in the imatinib arm, with 12.8%, 10.0% and 12.4% of patients were ≥ 65 years of age in the nilotinib 300 mg twice daily, nilotinib 400 mg twice daily and imatinib 400 mg once daily treatment arms, respectively. There were slightly more male than female patients (56.0%, 62.3% and 55.8%, in the nilotinib 300 mg twice daily, 400 mg twice daily and imatinib 400 mg once daily arm, respectively). More than 60% of all patients were Caucasian and 25% of all patients were Asian.

The primary data analysis time point was when all 846 patients completed 12 months of treatment (or discontinued earlier). Subsequent analyses reflect when patients completed 24, 36, 48, 60 and 72 months of treatment (or discontinued earlier). The median time on treatment was approximately 70 months in the nilotinib treatment groups and 64 months in the imatinib group. The median actual dose intensity was 593 mg/day for nilotinib 300 mg twice daily, 772 mg/day for nilotinib 400 mg twice daily and 400 mg/day for imatinib 400 mg once daily. This study is ongoing.

The primary efficacy endpoint was major molecular response (MMR) at 12 months. MMR was defined as $\leq 0.1\%$ BCR-ABL/ABL% by international scale (IS) measured by RQ-PCR, which corresponds to a ≥ 3 log reduction of BCR-ABL transcript from standardised baseline. The MMR rate at 12 months was statistically significantly higher for nilotinib 300 mg twice daily compared to imatinib 400 mg once daily (44.3% versus 22.3%, $p < 0.0001$). The rate of MMR at 12 months, was also statistically significantly higher for nilotinib 400 mg twice daily compared to imatinib 400 mg once daily (42.7% versus 22.3%, $p < 0.0001$).

The rates of MMR at 3, 6, 9 and 12 months were 8.9%, 33.0%, 43.3% and 44.3% for nilotinib 300 mg twice daily, 5.0%, 29.5%, 38.1% and 42.7% for nilotinib 400 mg twice daily and 0.7%, 12.0%, 18.0% and 22.3% for imatinib 400 mg once daily.

The MMR rate at 12, 24, 36, 48, 60 and 72 months is presented in Table 6.

Table 6 MMR rate

	Nilotinib 300 mg twice daily n=282 (%)	Nilotinib 400 mg twice daily n=281 (%)	Imatinib 400 mg once daily n=283 (%)
MMR at 12 months			
Response (95% CI)	44.3 ¹ (38.4; 50.3)	42.7 ¹ (36.8; 48.7)	22.3 (17.6; 27.6)
MMR at 24 months			
Response (95% CI)	61.7 ¹ (55.8; 67.4)	59.1 ¹ (53.1; 64.9)	37.5 (31.8; 43.4)
MMR at 36 months²			
Response (95% CI)	58.5 ¹ (52.5; 64.3)	57.3 ¹ (51.3; 63.2)	38.5 (32.8; 44.5)
MMR at 48 months³			
Response (95% CI)	59.9 ¹ (54.0; 65.7)	55.2 (49.1; 61.1)	43.8 (38.0; 49.8)
MMR at 60 months⁴			
Response (95% CI)	62.8 (56.8; 68.4)	61.2 (55.2; 66.9)	49.1 (43.2; 55.1)
MMR at 72 months⁵			
Response (95% CI)	52.5 (46.5; 58.4)	57.7 (51.6; 63.5)	41.7 (35.9; 47.7)

¹ Cochran-Mantel-Haenszel (CMH) test p-value for response rate (vs. imatinib 400 mg) <0.0001

² Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 199 (35.2%) of all patients were not evaluable for MMR at 36 months (87 in the nilotinib 300 mg twice daily group and 112 in the imatinib group) due to missing/unevaluable PCR assessments (n=17), atypical transcripts at baseline (n=7), or discontinuation prior to the 36-month time point (n=175).


³ Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 305 (36.1%) of all patients were not evaluable for MMR at 48 months (98 in the nilotinib 300 mg twice daily group, 88 in the nilotinib 400 mg twice daily group and 119 in the imatinib group) due to missing/unevaluable PCR assessments (n=18), atypical transcripts at baseline (n=8), or discontinuation prior to the 48-month time point (n=279).

⁴ Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 322 (38.1%) of all patients were not evaluable for MMR at 60 months (99 in the nilotinib 300 mg twice daily group, 93 in the nilotinib 400 mg twice daily group and 130 in the imatinib group) due to missing/unevaluable PCR assessments (n=9), atypical transcripts at baseline (n=8) or discontinuation prior to the 60-month time point (n=305).

⁵ Only patients who were in MMR at a specific time point are included as responders for that time point. A total of 395 (46.7%) of all patients were not evaluable for MMR at 72 months (130 in the nilotinib 300 mg twice daily group, 110 in the nilotinib 400 mg twice daily group and 155 in the imatinib group) due to missing/unevaluable PCR assessments (n=25), atypical transcripts at baseline (n=8) or discontinuation prior to the 72-month time point (n=362).

MMR rates by different time points (including patients who achieved MMR at or before those time points as responders) are presented in the cumulative incidence of MMR (see Figure 1).

Figure 1 Cumulative incidence of MMR



For all Sokal risk groups, the MMR rates at all time points remained consistently higher in the two nilotinib groups than in the imatinib group.

In a retrospective analysis, 91% (234/258) of patients on nilotinib 300 mg twice daily achieved BCR-ABL levels $\leq 10\%$ at 3 months of treatment compared to 67% (176/264) of patients on imatinib 400 mg once daily. Patients with BCR-ABL levels $\leq 10\%$ at 3 months of treatment show a greater overall survival at 72 months compared to those who did not achieve this molecular response level (94.5% vs. 77.1% respectively [p=0.0005]).

Based on the Kaplan-Meier analysis of time to first MMR the probability of achieving MMR at different time points was higher for both nilotinib at 300 mg and 400 mg twice daily compared to imatinib 400 mg once daily (HR=2.17 and stratified log-rank p<0.0001 between nilotinib 300 mg twice daily and imatinib 400 mg once daily, HR=1.88 and stratified log-rank p<0.0001 between nilotinib 400 mg twice daily and imatinib 400 mg once daily).

The proportion of patients who had a molecular response of $\leq 0.01\%$ and $\leq 0.0032\%$ by IS at different time points are presented in Table 7 and the proportion of patients who had a molecular response of $\leq 0.01\%$ and $\leq 0.0032\%$ by IS by different time points are presented in Figures 2 and 3. Molecular responses of $\leq 0.01\%$ and $\leq 0.0032\%$ by IS correspond to a ≥ 4 log reduction and ≥ 4.5 log reduction, respectively, of BCR-ABL transcripts from a standardised baseline.

Table 7 Proportions of patients who had molecular response of $\leq 0.01\%$ (4 log reduction) and $\leq 0.0032\%$ (4.5 log reduction)

	Nilotinib 300 mg twice daily n=282 (%)		Nilotinib 400 mg twice daily n=281 (%)		Imatinib 400 mg once daily n=283 (%)	
	$\leq 0.01\%$	$\leq 0.0032\%$	$\leq 0.01\%$	$\leq 0.0032\%$	$\leq 0.01\%$	$\leq 0.0032\%$
At 12 months	11.7	4.3	8.5	4.6	3.9	0.4
At 24 months	24.5	12.4	22.1	7.8	10.2	2.8
At 36 months	29.4	13.8	23.8	12.1	14.1	8.1
At 48 months	33.0	16.3	29.9	17.1	19.8	10.2
At 60 months	47.9	32.3	43.4	29.5	31.1	19.8
At 72 months	44.3	31.2	45.2	28.8	27.2	18.0

Figure 2 Cumulative incidence of molecular response of $\leq 0.01\%$ (4-log reduction)





Figure 3 Cumulative incidence of molecular response of $\leq 0.0032\%$ (4.5 log reduction)



Based on Kaplan-Meier estimates of the duration of first MMR, the proportions of patients who were maintaining response for 72 months among patients who achieved MMR were 92.5% (95% CI: 88.6-96.4%) in the nilotinib 300 mg twice daily group, 92.2% (95% CI: 88.5-95.9%) in the nilotinib 400 mg twice daily group and 88.0% (95% CI: 83.0-93.1%) in the imatinib 400 mg once daily group.

Complete cytogenetic response (CCyR) was defined as 0% Ph+ metaphases in the bone marrow based on a minimum of 20 metaphases evaluated. Best CCyR rate by 12 months (including patients who achieved CCyR at or before the 12 month time point as responders) was statistically higher for both nilotinib 300 mg and 400 mg twice daily compared to imatinib 400 mg once daily, see Table 8.

CCyR rate by 24 months (includes patients who achieved CCyR at or before the 24 month time point as responders) was statistically higher for both the nilotinib 300 mg twice daily and 400 mg twice daily groups compared to the imatinib 400 mg once daily group.

Table 8 Best CCyR rate

	Nilotinib 300 mg twice daily n=282 (%)	Nilotinib 400 mg twice daily n=281 (%)	Imatinib 400 mg once daily n=283 (%)
By 12 months			
Response (95% CI)	80.1 (75.0; 84.6)	77.9 (72.6; 82.6)	65.0 (59.2; 70.6)
No response	19.9	22.1	35.0
CMH test p-value for response rate (versus imatinib 400 mg once daily)	<0.0001	0.0005	
By 24 months			
Response (95% CI)	86.9 (82.4; 90.6)	84.7 (79.9; 88.7)	77.0 (71.7; 81.8)
No response	13.1	15.3	23.0
CMH test p-value for response rate (versus imatinib 400 mg once daily)	0.0018	0.0160	

Based on Kaplan-Meier estimates, the proportions of patients who were maintaining response for 72 months among patients who achieved CCyR were 99.1% (95% CI: 97.9-100%) in the nilotinib 300 mg twice daily group, 98.7% (95% CI: 97.1-100%) in the nilotinib 400 mg twice daily group and 97.0% (95% CI: 94.7-99.4%) in the imatinib 400 mg once daily group.

Progression to accelerated phase (AP) or blast crisis (BC) on treatment is defined as the time from the date of randomisation to the first documented disease progression to accelerated phase or blast crisis or CML-related death. Progression to accelerated phase or blast crisis on treatment was observed in a total of 17 patients: 2 patients on nilotinib 300 mg twice daily, 3 patients on nilotinib 400 mg twice daily and 12 patients on imatinib 400 mg once daily. The estimated rates of patients free from progression to accelerated phase or blast crisis at 72 months were 99.3%, 98.7% and 95.2%, respectively (HR=0.1599 and stratified log-rank p=0.0059 between nilotinib 300 mg twice daily and imatinib once daily, HR=0.2457 and stratified log-rank p=0.0185 between nilotinib 400 mg twice daily and imatinib once daily). No new events of progressions to AP/BC were reported on-treatment since the 2-year analysis.

Including clonal evolution as a criterion for progression, a total of 25 patients progressed to accelerated phase or blast crisis on treatment by the cut-off date (3 in the nilotinib 300 mg twice daily group, 5 in the nilotinib 400 mg twice daily group and 17 in the imatinib 400 mg once daily group). The estimated rates of patients free from progression to accelerated phase or blast crisis including clonal evolution at 72 months were 98.7%, 97.9% and 93.2%, respectively (HR=0.1626 and stratified log-rank p=0.0009 between nilotinib 300 mg twice daily and imatinib once daily, HR=0.2848 and stratified log-rank p=0.0085 between nilotinib 400 mg twice daily and imatinib once daily).

A total of 55 patients died during treatment or during the follow-up after discontinuation of treatment. (21 in the nilotinib 300 mg twice daily group, 11 in the nilotinib 400 mg twice daily group and 23 in the imatinib 400 mg once daily group). Twenty-six (26) of these 55 deaths were related to CML (6 in the nilotinib 300 mg twice daily group, 4 in the nilotinib 400 mg twice daily group and 16 in the imatinib 400 mg once daily group). The estimated rates of patients alive at 72 months were 91.6%, 95.8% and 91.4%, respectively (HR=0.8934 and stratified log-rank p=0.7085 between nilotinib 300 mg twice daily and imatinib, HR=0.4632 and stratified log-rank p=0.0314 between nilotinib 400 mg twice daily and imatinib). Considering only CML-related deaths as events, the estimated rates of overall survival at 72 months were 97.7%, 98.5% and 93.9%, respectively (HR=0.3694 and stratified log-rank p=0.0302 between nilotinib 300 mg twice daily and imatinib, HR=0.2433 and stratified log-rank p=0.0061 between nilotinib 400 mg twice daily and imatinib).

Treatment discontinuation in Ph+ CML patients in chronic phase who have been treated with nilotinib as first-line therapy and who have achieved a sustained deep molecular response

In an open-label, single-arm study, 215 adult patients with Ph+ CML in chronic phase treated with nilotinib in first-line for ≥ 2 years who achieved MR4.5 as measured with the MolecularMD MRDx BCR-ABL test were enrolled to continue nilotinib treatment for additional 52 weeks (nilotinib consolidation phase). 190 of 215 patients (88.4%) entered the TFR phase after achieving a sustained deep molecular response during the consolidation phase, defined by the following criteria:

- the 4 last quarterly assessments (taken every 12 weeks) were at least MR4 (BCR-ABL/ABL $\leq 0.01\%$ IS), and maintained for one year
- the last assessment being MR4.5 (BCR-ABL/ABL $\leq 0.0032\%$ IS)
- no more than two assessments falling between MR4 and MR4.5 ($0.0032\% \text{ IS} < \text{BCR-ABL/ABL} \leq 0.01\% \text{ IS}$).


The primary endpoint was the percentage of patients in MMR at 48 weeks after starting the TFR phase (considering any patient who required re-initiation of treatment as non-responder). Of the 190 patients who entered the TFR phase, 98 patients (51.6% [95% CI: 44.2, 58.9]) were in MMR at 48 weeks.

Eighty-eight patients (46.3%) discontinued the TFR phase due to loss of MMR, and 1 (0.5%), 1 (0.5%), and 3 patients (1.6%) due to death from unknown cause, physician decision and subject decision, respectively. Among these 88 patients, 86 patients restarted nilotinib treatment and 2 patients permanently discontinued the study. Eighty-five of these 86 patients (98.8%) regained MMR, (one patient discontinued study permanently due to subject decision) and 76 patients (88.4%) regained MR4.5 by the time of the cut-off date.

The Kaplan-Meier (KM) estimated median time on nilotinib treatment to regain MMR and MR4.5 was 7.9 weeks (95% CI: 5.1, 8.0) and 13.1 weeks (95% CI: 12.3, 15.7), respectively. The KM estimated MMR and MR4.5 rates at 24 weeks of re-initiation were 98.8% (95% CI: 94.2, 99.9) and 90.9% (95% CI: 83.2, 96.0), respectively.

The KM estimate of median treatment-free survival (TFS) has not yet been reached (Figure 4); 99 of 190 patients (52.1%) did not have a TFS event.

Figure 4 Kaplan-Meier estimate of treatment-free survival after start of TFR (full analysis set)



Paediatric population

The safety and efficacy of nilotinib in paediatric patients with Ph+ CML in chronic phase have been investigated in two studies. A total of 69 paediatric patients (from 2 to <18 years of age) with either newly diagnosed Ph+ CML in chronic phase (n=25) or imatinib/dasatinib resistant or imatinib-intolerant Ph+ CML in chronic phase (n=44) received nilotinib treatment at a dose of 230 mg/m² twice daily, rounded to the nearest 50 mg dose (to a maximum single dose of 400 mg).

In the pooled CML patient population, the median actual dose intensity was 435.5 mg/m²/day (range: 149 to 517 mg/m²/day), and the median relative dose intensity was 94.7% (range: 32 to 112%). Forty patients (58.0%) had relative dose intensity superior to 90%. The median time on treatment with nilotinib was 13.80 months (range: 0.7-30.9 months).

In the resistant or intolerant CML patients, the major molecular response (MMR; BCR-ABL/ABL ≤0.1% IS) rate was 40.9% (95% CI: 26.3, 56.8) at 12 cycles, with 18 patients being in MMR. In the newly diagnosed CML patients, the MMR rate was 60.0% (95% CI: 38.7, 78.9) at 12 cycles, with 15 patients achieving MMR. In resistant or intolerant CML patients, the cumulative MMR rate was 47.7% by cycle 12. In newly diagnosed CML patients, the cumulative MMR rate was 64.0% by cycle 12.

Among the 21 resistant or intolerant CML patients who were in MMR at any time on treatment, the median time to first MMR was 2.76 months (95% CI: 0.03, 5.55). For the 17 newly diagnosed CML patients who achieved MMR, the median time to first MMR was 5.55 months (95% CI: 5.52, 5.75).

Among resistant or intolerant CML patients, the percentage of patients who achieved BCR-ABL/ABL ≤0.01% IS (MR4.0) by the cut-off date was 11.4%, while 4.5% of the patients achieved BCR-ABL/ABL ≤0.0032% IS (MR4.5). Among newly diagnosed patients, the percentage of patients who achieved MR4.0 was 32%, while 28.0% achieved MR4.5.

None of the 21 resistant or intolerant CML patients who were in MMR on treatment, had confirmed loss of MMR. Among the 17 newly diagnosed CML patients who achieved MMR, one patient had confirmed loss of MMR (the patient lost CHR due to an increase in basophil count, however, did not progress to AP/BC).

One resistant or intolerant CML patient progressed to AP/BC after about 10 months on treatment.

No deaths were reported on treatment or after treatment discontinuation in both studies.

5.2 Pharmacokinetic properties

Absorption

Peak concentrations of nilotinib are reached 3 hours after oral administration. Nilotinib absorption following oral administration was approximately 30%. The absolute bioavailability of nilotinib has not been determined. As compared to an oral drink solution (pH of 1.2 to 1.3), relative bioavailability of nilotinib capsule is approximately 50%. In healthy volunteers, C_{max} and area under the serum concentration-time curve (AUC) of nilotinib are increased by 112% and 82%, respectively, compared to fasting conditions when Tasigna is given with food. Administration of Tasigna 30 minutes or 2 hours after food increased bioavailability of nilotinib by 29% or 15%, respectively (see sections 4.2, 4.4 and 4.5).

Nilotinib absorption (relative bioavailability) might be reduced by approximately 48% and 22% in patients with total gastrectomy and partial gastrectomy, respectively.

Distribution

The blood-to-plasma ratio of nilotinib is 0.71. Plasma protein binding is approximately 98% on the

basis of *in vitro* experiments.

Biotransformation

Main metabolic pathways identified in healthy subjects are oxidation and hydroxylation. Nilotinib is the main circulating component in the serum. None of the metabolites contribute significantly to the pharmacological activity of nilotinib. Nilotinib is primarily metabolised by CYP3A4, with possible minor contribution from CYP2C8.

Elimination

After a single dose of radiolabelled nilotinib in healthy subjects, more than 90% of the dose was eliminated within 7 days, mainly in faeces (94% of the dose). Unchanged nilotinib accounted for 69% of the dose.

The apparent elimination half-life estimated from the multiple-dose pharmacokinetics with daily dosing was approximately 17 hours. Inter-patient variability in nilotinib pharmacokinetics was moderate to high.

Linearity/non-linearity

Steady-state nilotinib exposure was dose-dependent, with less than dose-proportional increases in systemic exposure at dose levels higher than 400 mg given as once-daily dosing. Daily systemic exposure to nilotinib with 400 mg twice-daily dosing at steady state was 35% higher than with 800 mg once-daily dosing. Systemic exposure (AUC) of nilotinib at steady state at a dose level of 400 mg twice daily was approximately 13.4% higher than at a dose level of 300 mg twice daily. The average nilotinib trough and peak concentrations over 12 months were approximately 15.7% and 14.8% higher following 400 mg twice daily dosing compared to 300 mg twice daily. There was no relevant increase in exposure to nilotinib when the dose was increased from 400 mg twice daily to 600 mg twice daily.

Steady-state conditions were essentially achieved by day 8. An increase in serum exposure to nilotinib between the first dose and steady state was approximately 2-fold for daily dosing and 3.8-fold for twice-daily dosing.

Bioavailability/bioequivalence studies

Single-dose administration of 400 mg nilotinib, using 2 hard capsules of 200 mg whereby the content of each hard capsule was dispersed in one teaspoon of apple sauce, was shown to be bioequivalent with a single-dose administration of 2 intact hard capsules of 200 mg.

Paediatric population

Following administration of nilotinib in paediatric patients at 230 mg/m² twice daily, rounded to the nearest 50 mg dose (to a maximum single dose of 400 mg), steady-state exposure and clearance of nilotinib were found to be similar (within 2-fold) to adult patients treated with 400 mg twice daily. The pharmacokinetic exposure of nilotinib following a single or multiple doses appeared to be comparable between paediatric patients from 2 years to <10 years and from ≥10 years to <18 years.

5.3 Preclinical safety data

Nilotinib has been evaluated in safety pharmacology, repeated dose toxicity, genotoxicity, reproductive toxicity, phototoxicity and carcinogenicity (rats and mice) studies.

Nilotinib did not have effects on CNS or respiratory functions. *In vitro* cardiac safety studies demonstrated a preclinical signal for QT prolongation, based upon block of hERG currents and prolongation of the action potential duration in isolated rabbit hearts by nilotinib. No effects were seen in ECG measurements in dogs or monkeys treated for up to 39 weeks or in a special telemetry study in

dogs.

Repeated-dose toxicity studies in dogs of up to 4 weeks' duration and in cynomolgus monkeys of up to 9 months' duration revealed the liver as the primary target organ of toxicity of nilotinib. Alterations included increased alanine aminotransferase and alkaline phosphatase activity and histopathology findings (mainly sinusoidal cell or Kupffer cell hyperplasia/hypertrophy, bile duct hyperplasia and periportal fibrosis). In general the changes in clinical chemistry were fully reversible after a four-week recovery period and the histological alterations showed partial reversibility. Exposures at the lowest dose levels at which the liver effects were seen were lower than the exposure in humans at a dose of 800 mg/day. Only minor liver alterations were seen in mice or rats treated for up to 26 weeks. Mainly reversible increases in cholesterol levels were seen in rats, dogs and monkeys.

Genotoxicity studies in bacterial *in vitro* systems and in mammalian *in vitro* and *in vivo* systems with and without metabolic activation did not reveal any evidence for a mutagenic potential of nilotinib.

In the 2-year rat carcinogenicity study, the major target organ for non-neoplastic lesions was the uterus (dilatation, vascular ectasia, endothelial cell hyperplasia, inflammation and/or epithelial hyperplasia). There was no evidence of carcinogenicity upon administration of nilotinib at 5, 15 and 40 mg/kg/day. Exposures (in terms of AUC) at the highest dose level represented approximately 2x to 3x human daily steady-state exposure (based on AUC) to nilotinib at the dose of 800 mg/day.

In the 26-week Tg.rasH2 mouse carcinogenicity study, in which nilotinib was administered at 30, 100 and 300 mg/kg/day, skin papillomas/carcinomas were detected at 300 mg/kg, representing approximately 30 to 40 times (based on AUC) the human exposure at the maximum approved dose of 800 mg/day (administered as 400 mg twice daily). The No-Observed-Effect-Level for the skin neoplastic lesions was 100 mg/kg/day, representing approximately 10 to 20 times the human exposure at the maximum approved dose of 800 mg/day (administered as 400 mg twice daily). The major target organs for non-neoplastic lesions were the skin (epidermal hyperplasia), the growing teeth (degeneration/atrophy of the enamel organ of upper incisors and inflammation of the gingiva/odontogenic epithelium of incisors) and the thymus (increased incidence and/or severity of decreased lymphocytes).

Nilotinib did not induce teratogenicity, but did show embryo- and foetotoxicity at doses that also showed maternal toxicity. Increased post-implantation loss was observed in both the fertility study, which involved treatment of both males and females, and the embryotoxicity study, which involved treatment of females. Embryo-lethality and foetal effects (mainly decreased foetal weights, premature fusion of the facial bones (fused maxilla/zygomatic) visceral and skeletal variations) in rats and increased resorption of foetuses and skeletal variations in rabbits were present in the embryotoxicity studies. In a pre- and postnatal development study in rats, maternal exposure to nilotinib caused reduced pup body weight with associated changes in physical development parameters as well as reduced mating and fertility indices in the offspring. Exposure to nilotinib in females at No-Observed-Adverse-Effect-Levels was generally less or equal to that in humans at 800 mg/day.

In a juvenile development study, nilotinib was administered via oral gavage to juvenile rats from the first week post partum through young adult (day 70 post partum) at doses of 2, 6 and 20 mg/kg/day. Besides standard study parameters, evaluations of developmental landmarks, CNS effects, mating and fertility were performed. Based on a reduction in body weight in both genders and a delayed preputial separation in males (which may be associated with the reduction in weight), the No-Observed-Effect-Level in juvenile rats was considered to be 6 mg/kg/day. The juvenile animals did not exert increased sensitivity to nilotinib relative to adults. In addition, the toxicity profile in juvenile rats was comparable to that observed in adult rats.

No effects on sperm count/motility or on fertility were noted in male and female rats up to the highest tested dose, approximately 5 times the recommended dosage for humans.

Nilotinib was shown to absorb light in the UV-B and UV-A range, is distributed into the skin and showed a phototoxic potential *in vitro*, but no effects have been observed *in vivo*. Therefore the risk

that nilotinib causes photosensitisation in patients is considered very low.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content

Lactose monohydrate
Crospovidone Type A
Poloxamer 188
Colloidal anhydrous silica
Magnesium stearate

Capsule shell

Gelatin
Titanium dioxide (E171)
Red iron oxide (E172)
Yellow iron oxide (E172)

Printing ink

Shellac
Black iron oxide (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Do not store above 30°C.

Store in the original package in order to protect from moisture.

6.5 Nature and contents of container

PVC/PVDC/Alu blisters.

Tasigna is available in the following pack sizes:

- Unit packs containing 28 hard capsules (7 daily blisters, each containing 4 hard capsules) or 40 hard capsules (5 blisters, each containing 8 hard capsules).
- Multipacks containing 112 (4 packs of 28) hard capsules, 120 (3 packs of 40) hard capsules or 392 (14 packs of 28) hard capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/005-006
EU/1/07/422/009-010
EU/1/07/422/013

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 19 November 2007
Date of latest renewal: 19 November 2012

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

ANNEX II

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

Novartis Pharma GmbH
Roonstraße 25
D-90429 Nuremberg
Germany

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (See Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

• Periodic Safety Update Reports

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

• Risk Management Plan (RMP)

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.
- **Additional risk minimisation measures**

The MAH shall ensure that prior to launch, all doctors who intend to prescribe the medicinal product, and all pharmacists who may dispense the medicinal product, are provided with a healthcare professional information pack containing the following:

- Educational brochure
- Summary of Product Characteristics (SPC) and Package Leaflet and Labelling

Key elements to be included in the educational brochure

- Brief background on Tasigna, its authorised indication and posology
- Information on the cardiac risks associated with the use of Tasigna
 - That Tasigna can cause prolongation of the QT interval and that Tasigna should be used with caution in patients who have or who are at significant risk of developing prolongation of QTc. Concomitant use of Tasigna with anti-arrhythmics or other

- medicinal products that may prolong the QT interval should be undertaken with caution
 - Caution in prescribing to patients with a history of or risk factors for coronary heart disease
 - That Tasigna may cause fluid retention, cardiac failure and pulmonary oedema
- That Tasigna is metabolised by CYP3A4 and that strong inhibitors or inducers of this enzyme may significantly affect exposure to Tasigna.
 - That inhibitors may increase the potential for adverse drug reactions in particular QT interval prolongation.
 - To warn patients about OTC medicines in particular St John's Wort
- The need to inform patients about the effects of food on Tasigna
 - Not to eat within two hours before and one hour after taking Tasigna
 - The need to avoid foods such as grapefruit juice which inhibit CYP3A4 enzymes
- **Obligation to conduct post-authorisation measures**

The MAH shall complete, within the stated timeframe, the below measures:

Description	Due date
Post authorisation efficacy study (PAES): In order to investigate the efficacy of nilotinib in paediatric patients with Ph+ CML-CP resistant or intolerant to either imatinib or dasatinib and in newly diagnosed patients, the MAH should submit the final results of the phase II CAMN107A2203 study.	Final CSR: April 2021

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 50 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 50 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

120 (3 packs of 40) hard capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/015 120 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Tasigna 50 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

INTERMEDIATE CARTON

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 50 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 50 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

40 hard capsules. Not to be sold separately.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/015 120 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Tasigna 50 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTERS

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 50 mg hard capsules
nilotinib

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON OF UNIT PACK

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 150 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 150 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

28 hard capsules

40 hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/005	28 hard capsules
EU/1/07/422/009	40 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Tasigna 150 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTERS

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 150 mg hard capsules
nilotinib

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON OF MULTIPACK (INCLUDING BLUE BOX)

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 150 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 150 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

Multipack: 112 (4 packs of 28) hard capsules.
Multipack: 120 (3 packs of 40) hard capsules.
Multipack: 392 (14 packs of 28) hard capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/006	112 hard capsules
EU/1/07/422/010	120 hard capsules
EU/1/07/422/013	392 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Tasigna 150 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

INTERMEDIATE CARTON OF MULTIPACK (WITHOUT BLUE BOX)

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 150 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 150 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

28 hard capsules. Component of a multipack. Not to be sold separately.

40 hard capsules. Component of a multipack. Not to be sold separately.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/006	112 hard capsules
EU/1/07/422/010	120 hard capsules
EU/1/07/422/013	392 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Tasigna 150 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON OF UNIT PACK (WALLET)
CARTON OF UNIT PACK (CARTON)

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 200 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 200 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

28 hard capsules

40 hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/001	PVC/PVDC/Alu [in wallet] 28 hard capsules
EU/1/07/422/002	PA/Alu/PVC/Alu [in wallet] 28 hard capsules
EU/1/07/422/007	PVC/PVDC/Alu [in carton] 28 hard capsules
EU/1/07/422/011	PVC/PVDC/Alu [in carton] 40 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Tasigna 200 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTERS

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 200 mg hard capsules
nilotinib

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON OF MULTIPACK (WALLET) (INCLUDING BLUE BOX)
CARTON OF MULTIPACK (CARTON) (INCLUDING BLUE BOX)

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 200 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 200 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

Multipack: 112 (4 wallets of 28) hard capsules.
Multipack: 112 (4 packs of 28) hard capsules.
Multipack: 120 (3 packs of 40) hard capsules.
Multipack: 392 (14 packs of 28) hard capsules.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/003	PVC/PVDC/Alu [in wallet] 112 hard capsules
EU/1/07/422/004	PA/Alu/PVC/Alu [in wallet] 112 hard capsules
EU/1/07/422/008	PVC/PVDC/Alu [in carton] 112 hard capsules
EU/1/07/422/012	PVC/PVDC/Alu [in carton] 120 hard capsules
EU/1/07/422/014	PVC/PVDC/Alu [in carton] 392 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Tasigna 200 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

INTERMEDIATE WALLET OF MULTIPACK (WITHOUT BLUE BOX)
INTERMEDIATE CARTON OF MULTIPACK (WITHOUT BLUE BOX)

1. NAME OF THE MEDICINAL PRODUCT

Tasigna 200 mg hard capsules
nilotinib

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One hard capsule contains 200 mg nilotinib (as hydrochloride monohydrate).

3. LIST OF EXCIPIENTS

Contains lactose – see the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

28 hard capsules. Component of a multipack comprising 4 wallets. Not to be sold separately.
28 hard capsules. Component of a multipack comprising 4 cartons. Not to be sold separately.
40 hard capsules. Component of a multipack comprising 3 cartons. Not to be sold separately.
28 hard capsules. Component of a multipack comprising 14 cartons. Not to be sold separately.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/07/422/003	PVC/PVDC/Alu [in wallet] 112 hard capsules
EU/1/07/422/004	PA/Alu/PVC/Alu [in wallet] 112 hard capsules
EU/1/07/422/008	PVC/PVDC/Alu [in carton] 112 hard capsules
EU/1/07/422/012	PVC/PVDC/Alu [in carton] 120 hard capsules
EU/1/07/422/014	PVC/PVDC/Alu [in carton] 392 hard capsules

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Tasigna 200 mg

17. UNIQUE IDENTIFIER – 2D BARCODE**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

B. PACKAGE LEAFLET

Package leaflet: Information for the user

Tasigna 50 mg hard capsules nilotinib

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Tasigna is and what it is used for
2. What you need to know before you take Tasigna
3. How to take Tasigna
4. Possible side effects
5. How to store Tasigna
6. Contents of the pack and other information

1. What Tasigna is and what it is used for

What Tasigna is

Tasigna is a medicine containing an active substance called nilotinib.

What Tasigna is used for

Tasigna is used to treat a type of leukaemia called Philadelphia chromosome positive chronic myeloid leukaemia (Ph-positive CML). CML is a cancer of the blood which makes the body produce too many abnormal white blood cells.

Tasigna is used in adult and paediatric patients with newly diagnosed CML or in patients with CML who are no longer benefiting from previous treatment including imatinib. It is also used in adult and paediatric patients who experienced serious side effects with previous treatment and are not able to continue taking it.

How Tasigna works

In patients with CML, a change in DNA (genetic material) triggers a signal that tells the body to produce abnormal white blood cells. Tasigna blocks this signal, and thus stops the production of these cells.

Monitoring during Tasigna treatment

Regular tests, including blood tests, will be performed during treatment. These tests will monitor:

- the amount of blood cells (white blood cells, red blood cells and platelets) in the body to see how Tasigna is tolerated.
- pancreas and liver function in the body to see how Tasigna is tolerated.
- the electrolytes in the body (potassium, magnesium). These are important in the functioning of the heart.
- the level of sugar and fats in the blood.

The heart rate will also be checked using a machine that measures electrical activity of the heart (a test called an "ECG").

Your doctor will regularly evaluate your treatment and decide whether you should continue to take Tasigna. If you are told to discontinue this medicine, your doctor will continue to monitor your CML and may tell you to re-start Tasigna if your condition indicates that this is necessary.

If you have any questions about how Tasigna works or why it has been prescribed for you or your child, ask your doctor.

2. What you need to know before you take Tasigna

Follow all the doctor's instructions carefully. They may differ from the general information contained in this leaflet.

Do not take Tasigna

- if you are allergic to nilotinib or any of the other ingredients of this medicine (listed in section 6).

If you think you may be allergic, tell your doctor **before taking Tasigna**.

Warnings and precautions

Talk to your doctor or pharmacist before taking Tasigna:

- if you have suffered prior cardiovascular events such as a heart attack, chest pain (angina), problems with the blood supply to your brain (stroke) or problems with the blood flow to your leg (claudication) or if you have risk factors for cardiovascular disease such as high blood pressure (hypertension), diabetes or problems with the level of fats in your blood (lipid disorders).
- if you have a **heart disorder**, such as an abnormal electrical signal called "prolongation of the QT interval".
- if you are being **treated with medicines** that affect the heart beat (anti-arrhythmics) or the liver (see **Other medicines and Tasigna**).
- if you suffer from lack of potassium or magnesium.
- if you have a liver or pancreas disorder.
- if you have symptoms such as easy bruising, feeling tired or short of breath or have experienced repeated infections.
- if you have had a surgical procedure involving the removal of the entire stomach (total gastrectomy).
- if you have ever had or might now have a hepatitis B infection. This is because Tasigna could cause hepatitis B to become active again, which can be fatal in some cases. Patients will be carefully checked by their doctor for signs of this infection before treatment is started.

If any of these apply to you or your child, tell your doctor.

During treatment with Tasigna

- if you faint (loss of consciousness) or have an irregular heart beat while taking this medicine, **tell your doctor immediately** as this may be a sign of a serious heart condition. Prolongation of the QT interval or an irregular heart beat may lead to sudden death. Uncommon cases of sudden death have been reported in patients taking Tasigna.
- if you have sudden heart palpitations, severe muscle weakness or paralysis, seizures or sudden changes in your thinking or level of alertness, **tell your doctor immediately** as this may be a sign of a fast breakdown of cancer cells called tumour lysis syndrome. Rare cases of tumour lysis syndrome have been reported in patients treated with Tasigna.
- if you develop chest pain or discomfort, numbness or weakness, problems with walking or with your speech, pain, discolouration or a cool feeling in a limb, **tell your doctor immediately** as this may be a sign of a cardiovascular event. Serious cardiovascular events including problems with the blood flow to the leg (peripheral arterial occlusive disease), ischaemic heart disease and problems with the blood supply to the brain (ischaemic cerebrovascular disease) have been reported in patients taking Tasigna. Your doctor should assess the level of fats (lipids) and sugar in your blood before initiating treatment with Tasigna and during treatment.
- if you develop swelling of the feet or hands, generalised swelling or rapid weight gain tell your

doctor as these may be signs of severe fluid retention. Uncommon cases of severe fluid retention have been reported in patients treated with Tasigna.

If you are the parent of a child who is being treated with Tasigna, tell the doctor if any of the above conditions apply to your child.

Children and adolescents

Tasigna is a treatment for children and adolescents with CML. There is no experience with the use of this medicine in children below 2 years of age. There is no experience with the use of Tasigna in newly diagnosed children below 10 years of age and limited experience in patients below 6 years of age who are no longer benefiting from previous treatment for CML. The long-term effects of treating children with Tasigna for long periods of time are not known.

Other medicines and Tasigna

Tasigna may interfere with some other medicines.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes in particular:

- anti-arrhythmics – used to treat irregular heart beat;
- chloroquine, halofantrine, clarithromycin, haloperidol, methadone, moxifloxacin - medicines that may have an unwanted effect on the function of the heart;
- ketoconazole, itraconazole, voriconazole, clarithromycin, telithromycin – used to treat infections;
- ritonavir – a medicine from the class “ antiproteases” used to treat HIV;
- carbamazepine, phenobarbital, phenytoin – used to treat epilepsy;
- rifampicin – used to treat tuberculosis;
- St. John’s Wort – a herbal product used to treat depression and other conditions (also known as *Hypericum perforatum*);
- midazolam – used to relieve anxiety before surgery;
- alfentanil and fentanyl – used to treat pain and as a sedative before or during surgery or medical procedures;
- cyclosporine, sirolimus and tacrolimus – medicines that suppress the “self-defence” ability of the body and fight infections and are commonly used to prevent the rejection of transplanted organs such as the liver, heart and kidney;
- dihydroergotamine and ergotamine – used to treat dementia;
- lovastatin, simvastatin – used to treat high level of fats in blood;
- warfarin – used to treat blood coagulation disorders (such as blood clots or thromboses);
- astemizole, terfenadine, cisapride, pimozone, quinidine, bepridil or ergot alkaloids (ergotamine, dihydroergotamine).

These medicines should be avoided during your treatment with Tasigna. If you are taking any of these, your doctor might prescribe other alternative medicines.

In addition, tell your doctor or pharmacist before taking Tasigna if you are taking any antacids, which are medicines against heartburn. These medicines need to be taken separately from Tasigna:

- H2 blockers, which decrease the production of acid in the stomach. H2 blockers should be taken approximately 10 hours before and approximately 2 hours after you take Tasigna;
- antacids such as those containing aluminium hydroxide, magnesium hydroxide and simethicone, which neutralise high acidity in the stomach. These antacids should be taken approximately 2 hours before or approximately 2 hours after you take Tasigna.

You should also tell your doctor **if you are already taking Tasigna** and you are prescribed a new medicine that you have not taken previously during Tasigna treatment.

Tasigna with food and drink

Do not take Tasigna with food. Food may enhance the absorption of Tasigna and therefore increase the amount of Tasigna in the blood, possibly to a harmful level. Do not drink grapefruit juice or eat grapefruit. It may increase the amount of Tasigna in the blood, possibly to a harmful level.

Pregnancy and breast-feeding

- **Tasigna is not recommended during pregnancy** unless clearly necessary. If you are pregnant or think that you may be, tell your doctor who will discuss with you whether you can take this medicine during your pregnancy.
- **Women who might get pregnant** are advised to use highly effective contraception during treatment and for up to two weeks after ending treatment.
- **Breast-feeding is not recommended** during treatment with Tasigna. Tell your doctor if you are breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

If you experience side effects (such as dizziness or visual disorders) with a potential impact on the ability to safely drive or use any tools or machines after taking this medicine, you should refrain from these activities until the effect has disappeared.

Tasigna contains lactose

This medicine contains lactose (also known as milk sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Tasigna

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much Tasigna to take

Use in adults

- **Patients newly diagnosed with CML:** The recommended dose is 600 mg per day. This dose is achieved by taking two hard capsules of 150 mg twice a day.
- **Patients who are no longer benefiting from previous treatment for CML:** The recommended dose is 800 mg per day. This dose is achieved by taking two hard capsules of 200 mg twice a day.

Use in children and adolescents

- The dose given to your child will depend on your child's body weight and height. The doctor will calculate the correct dose to use and tell you how many capsules of Tasigna to give to your child. The total daily dose you give to your child must not exceed 800 mg.

Your doctor may prescribe a lower dose depending on how you respond to treatment.

Older people (age 65 years and over)

Tasigna can be used by people aged 65 years and over at the same dose as for other adults.

When to take Tasigna

Take the hard capsules:

- twice a day (approximately every 12 hours);
- at least 2 hours after any food;
- then wait 1 hour before eating again.

If you have questions about when to take this medicine, talk to your doctor or pharmacist. Taking Tasigna at the same time each day will help you remember when to take your hard capsules.

How to take Tasigna

- Swallow the hard capsules whole with water.
- Do not take any food together with the hard capsules.

- Do not open the hard capsules unless you are unable to swallow them. If so, you may sprinkle the content of each hard capsule in **one** teaspoon of apple sauce and take it immediately. Do not use more than one teaspoon of apple sauce for each hard capsule and do not use any food other than apple sauce.

How long to take Tasigna

Continue taking Tasigna every day for as long as your doctor tells you. This is a long-term treatment. Your doctor will regularly monitor your condition to check that the treatment is having the desired effect.

Your doctor may consider discontinuing your treatment with Tasigna based on specific criteria.

If you have questions about how long to take Tasigna, talk to your doctor.

If you take more Tasigna than you should

If you have taken more Tasigna than you should have, or if someone else accidentally takes your hard capsules, contact a doctor or hospital for advice straight away. Show them the pack of hard capsules and this package leaflet. Medical treatment may be necessary.

If you forget to take Tasigna

If you miss a dose, take your next dose as scheduled. Do not take a double dose to make up for a forgotten hard capsule.

If you stop taking Tasigna

Do not stop taking this medicine unless your doctor tells you to do so. Stopping Tasigna without your doctor's recommendation places you at risk for worsening of your disease which could have life-threatening consequences. Be sure to discuss with your doctor, nurse, and/or pharmacist if you are considering stopping Tasigna.

If your doctor recommends that you discontinue treatment with Tasigna

Your doctor will regularly evaluate your treatment with a specific diagnostic test and decide whether you should continue to take this medicine. If you are told to discontinue Tasigna, your doctor will continue to carefully monitor your CML before, during and after you have discontinued Tasigna and may tell you to re-start Tasigna if your condition indicates that this is necessary.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the side effects are mild to moderate and will generally disappear after a few days to a few weeks of treatment.

Some side effects could be serious.

These side effects are common (may affect up to 1 in 10 people), uncommon (may affect up to 1 in 100 people) or have been reported with frequency not known (cannot be estimated from the available data).

- rapid weight gain, swelling of hands, ankles, feet or face (signs of water retention)
- chest pain, high blood pressure, irregular heart rhythm, blue discolouration of the lips, tongue or skin (signs of heart disorders)
- difficulty breathing, cough, wheezing with or without fever, swelling of the feet or legs (signs of lung disorders)
- fever, easy bruising, frequent infections (signs of blood disorders)
- weakness or paralysis of the limbs or face, difficulty speaking, severe headache, seeing, feeling or hearing things that are not there (signs of nervous system disorders)
- thirst, dry skin, irritability, dark urine, decreased urine output (signs of kidney disorders)
- blurred vision, loss of vision, blood in eye (signs of eye disorders)
- swelling and pain in one part of the body (signs of clotting within a vein)

- abdominal pain, nausea, vomiting of blood, black stools, constipation, swollen abdomen (signs of gastrointestinal disorders)
- severe upper abdominal pain (sign of pancreatitis)
- yellow skin and eyes, nausea, loss of appetite, dark-coloured urine (signs of liver disorders)
- rash, painful red lumps, pain in joints and muscles (signs of skin disorders)
- excessive thirst, high urine output, increased appetite with weight loss, tiredness (signs of high level of sugar in the blood)
- fast heartbeat, bulging eyes, weight loss, swelling at the front of the neck (signs of overactive thyroid gland)
- nausea, shortness of breath, irregular heartbeat, clouding of urine, tiredness and/or joint discomfort associated with abnormal results of blood tests (such as high levels of potassium, uric acid and phosphorous and low levels of calcium)
- pain, discomfort, weakness or cramping in the leg muscles, which may be due to decreased blood flow, ulcers on the legs or arms that heal slowly or not at all and noticeable changes in colour (blueness or paleness) or temperature (coolness) of the legs or arms, as these symptoms could be signs of artery blockage in the affected limb (leg or arm) and digits (toes and fingers)
- recurrence (reactivation) of hepatitis B infection when you have had hepatitis B in the past (a liver infection).

If you get any of these, **tell your doctor straight away.**

Some side effects are very common (may affect more than 1 in 10 people)

- diarrhoea
- headache
- tiredness
- muscle pain
- itching, rash
- nausea
- vomiting
- hair loss
- high blood level of bilirubin (liver function)
- high blood level of lipase (pancreas function)
- musculoskeletal pain, muscle pain, pain in extremity, pain in joints, bone pain and spinal pain upon discontinuing treatment with Tasigna

If any of these affects you severely, tell your doctor.

Some side effects are common (may affect up to 1 in 10 people)

- abdominal discomfort, stomach discomfort after meals, flatulence, swelling or bloating of the abdomen
- bone pain, pain in joints, muscle spasms
- pain including back pain, neck pain and pain in extremity, pain or discomfort in the side of the body
- eye irritation, swelling, discharge, itching or redness, dry eye (signs of eye disorders)
- skin reddening, dry skin, acne, wart, decreased skin sensitivity, hives
- loss of appetite, disturbed sense of taste, weight decrease or increase
- insomnia, depression, anxiety
- night sweats, excessive sweating, hot flushes
- dizziness, generally feeling unwell, spinning sensation
- tingling or numbness
- voice disorder
- nose bleed
- frequent urine output
- palpitations (sensation of rapid heart beat)

If any of these affects you severely, tell your doctor.

Some side effects are uncommon (may affect up to 1 in 100 people)

- increased skin sensitivity, skin pain
- swelling of the eyelids

- dry mouth, sore throat, mouth sores
- heartburn
- breast pain
- increased appetite
- attention disorder
- difficulty and pain when urinating, exaggerated sense of needing to urinate
- inability to achieve or maintain an erection
- breast enlargement in men
- flu-like symptoms, muscle weakness
- trembling
- decreased sharpness of vision
- severe headache often accompanied by nausea, vomiting and sensitivity to light
- visual disturbances
- oral or vaginal thrush
- muscle and joint stiffness
- unconsciousness
- weight gain
- feeling body temperature change (including feeling hot, feeling cold)
- thickened patches of red/silver skin (signs of psoriasis)
- sensitive teeth

If any of these affects you severely, tell your doctor.

The following other side effects have been reported with frequency not known (cannot be estimated from the available data):

- confusion, disorientation, memory loss, disturbed mood, lack of energy
- bacterial infection of the skin
- blister, skin cyst, oily skin, thinning of the skin, dark patches of skin, skin discolouration
- bleeding, tender or enlarged gums
- runny or stuffy nose, sneezing
- reddening and/or swelling and possibly peeling on the palms and soles (so called hand-foot syndrome)
- increased sensitivity of the eyes or the skin to light
- eye pain or redness, pain, itching of the eyelids
- difficulty hearing, ear pain, noises (ringing) in the ears
- painful and swollen joints (gout)
- blood in urine, abnormal urine colour, urinary incontinence
- haemorrhoids
- feeling of hardening in the breasts, heavy periods, nipple swelling
- symptoms of restless legs syndrome (an irresistible urge to move a part of the body, usually the leg, accompanied by uncomfortable sensations)

If any of these affects you severely, tell your doctor.

During Tasigna treatment, you may also have some abnormal blood test results such as low level of blood cells (white cells, red cells, platelets), high blood level of lipase or amylase (pancreas function), high blood level of bilirubin (liver function) or high blood level of creatinine (kidney function), low or high blood level of insulin (a hormone regulating blood sugar level), low or high level of sugar, or high level of fats in the blood.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Tasigna

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Store in the original package in order to protect from moisture.
- Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Tasigna contains

- The active substance is nilotinib. Each hard capsule contains 50 mg nilotinib (as hydrochloride monohydrate).
- The other ingredients are lactose monohydrate, crospovidone type A, poloxamer 188, colloidal anhydrous silica, magnesium stearate. The hard capsule shell is composed of gelatin, titanium dioxide (E171), red iron oxide (E172), yellow iron oxide (E172) and, shellac, black iron oxide (E172), propylene glycol and ammonium hydroxide for stamping of the imprint.

What Tasigna looks like and contents of the pack

Tasigna is supplied as hard capsules. The hard capsules are red/light yellow. A black imprint is stamped on each hard capsule (“NVR/ABL”).

Tasigna is available in a pack containing 120 hard capsules (3 packs of 40 hard capsules).

Marketing Authorisation Holder

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

Manufacturer

Novartis Pharma GmbH
Roonstraße 25
D-90429 Nuremberg
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Novartis Pharma N.V.
Tél/Tel: +32 2 246 16 11

Lietuva

Novartis Pharma Services Inc.
Tel: +370 5 269 16 50

България

Novartis Bulgaria EOOD
Тел.: +359 2 489 98 28

Luxembourg/Luxemburg

Novartis Pharma N.V.
Tél/Tel: +32 2 246 16 11

Česká republika

Novartis s.r.o.
Tel: +420 225 775 111

Danmark

Novartis Healthcare A/S
Tlf: +45 39 16 84 00

Deutschland

Novartis Pharma GmbH
Tel: +49 911 273 0

Eesti

Novartis Pharma Services Inc.
Tel: +372 66 30 810

Ελλάδα

Novartis (Hellas) A.E.B.E.
Τηλ: +30 210 281 17 12

España

Novartis Farmacéutica, S.A.
Tel: +34 93 306 42 00

France

Novartis Pharma S.A.S.
Tél: +33 1 55 47 66 00

Hrvatska

Novartis Hrvatska d.o.o.
Tel. +385 1 6274 220

Ireland

Novartis Ireland Limited
Tel: +353 1 260 12 55

Ísland

Vistor hf.
Sími: +354 535 7000

Italia

Novartis Farma S.p.A.
Tel: +39 02 96 54 1

Κύπρος

Novartis Pharma Services Inc.
Τηλ: +357 22 690 690

Latvija

SIA "Novartis Baltics"
Tel: +371 67 887 070

Magyarország

Novartis Hungária Kft.
Tel.: +36 1 457 65 00

Malta

Novartis Pharma Services Inc.
Tel: +356 2122 2872

Nederland

Novartis Pharma B.V.
Tel: +31 26 37 82 555

Norge

Novartis Norge AS
Tlf: +47 23 05 20 00

Österreich

Novartis Pharma GmbH
Tel: +43 1 86 6570

Polska

Novartis Poland Sp. z o.o.
Tel.: +48 22 375 4888

Portugal

Novartis Farma - Produtos Farmacêuticos, S.A.
Tel: +351 21 000 8600

România

Novartis Pharma Services Romania SRL
Tel: +40 21 31299 01

Slovenija

Novartis Pharma Services Inc.
Tel: +386 1 300 75 50

Slovenská republika

Novartis Slovakia s.r.o.
Tel: +421 2 5542 5439

Suomi/Finland

Novartis Finland Oy
Puh/Tel: +358 (0)10 6133 200

Sverige

Novartis Sverige AB
Tel: +46 8 732 32 00

United Kingdom

Novartis Pharmaceuticals UK Ltd.
Tel: +44 1276 698370

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu/>. There are also links to other websites about rare diseases and treatments.

Package leaflet: Information for the user

Tasigna 150 mg hard capsules nilotinib

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Tasigna is and what it is used for
2. What you need to know before you take Tasigna
3. How to take Tasigna
4. Possible side effects
5. How to store Tasigna
6. Contents of the pack and other information

1. What Tasigna is and what it is used for

What Tasigna is

Tasigna is a medicine containing an active substance called nilotinib.

What Tasigna is used for

Tasigna is used to treat a type of leukaemia called Philadelphia chromosome positive chronic myeloid leukaemia (Ph-positive CML). CML is a cancer of the blood which makes the body produce too many abnormal white blood cells.

Tasigna is used in adult and paediatric patients with newly diagnosed CML. It is also used in paediatric patients with CML who are no longer benefiting from previous treatment including imatinib or who experienced serious side effects with previous treatment and are not able to continue taking it.

How Tasigna works

In patients with CML, a change in DNA (genetic material) triggers a signal that tells the body to produce abnormal white blood cells. Tasigna blocks this signal, and thus stops the production of these cells.

Monitoring during Tasigna treatment

Regular tests, including blood tests, will be performed during treatment. These tests will monitor:

- the amount of blood cells (white blood cells, red blood cells and platelets) in the body to see how Tasigna is tolerated.
- pancreas and liver function in the body to see how Tasigna is tolerated.
- the electrolytes in the body (potassium, magnesium). These are important in the functioning of the heart.
- the level of sugar and fats in the blood.

The heart rate will also be checked using a machine that measures electrical activity of the heart (a test called an "ECG").

Your doctor will regularly evaluate your treatment and decide whether you should continue to take Tasigna. If you are told to discontinue this medicine, your doctor will continue to monitor your CML and may tell you to re-start Tasigna if your condition indicates that this is necessary.

If you have any questions about how Tasigna works or why it has been prescribed for you or your child, ask your doctor.

2. What you need to know before you take Tasigna

Follow all the doctor's instructions carefully. They may differ from the general information contained in this leaflet.

Do not take Tasigna

- if you are allergic to nilotinib or any of the other ingredients of this medicine (listed in section 6).

If you think you may be allergic, tell your doctor **before taking Tasigna**.

Warnings and precautions

Talk to your doctor or pharmacist before taking Tasigna:

- if you have suffered prior cardiovascular events such as a heart attack, chest pain (angina), problems with the blood supply to your brain (stroke) or problems with the blood flow to your leg (claudication) or if you have risk factors for cardiovascular disease such as high blood pressure (hypertension), diabetes or problems with the level of fats in your blood (lipid disorders).
- if you have a **heart disorder**, such as an abnormal electrical signal called "prolongation of the QT interval".
- if you are being **treated with medicines** that affect the heart beat (anti-arrhythmics) or the liver (see **Other medicines and Tasigna**).
- if you suffer from lack of potassium or magnesium.
- if you have a liver or pancreas disorder.
- if you have symptoms such as easy bruising, feeling tired or short of breath or have experienced repeated infections.
- if you have had a surgical procedure involving the removal of the entire stomach (total gastrectomy).
- if you have ever had or might now have a hepatitis B infection. This is because Tasigna could cause hepatitis B to become active again, which can be fatal in some cases. Patients will be carefully checked by their doctor for signs of this infection before treatment is started.

If any of these apply to you or your child, tell your doctor.

During treatment with Tasigna

- if you faint (loss of consciousness) or have an irregular heart beat while taking this medicine, **tell your doctor immediately** as this may be a sign of a serious heart condition. Prolongation of the QT interval or an irregular heart beat may lead to sudden death. Uncommon cases of sudden death have been reported in patients taking Tasigna.
- if you have sudden heart palpitations, severe muscle weakness or paralysis, seizures or sudden changes in your thinking or level of alertness, **tell your doctor immediately** as this may be a sign of a fast breakdown of cancer cells called tumour lysis syndrome. Rare cases of tumour lysis syndrome have been reported in patients treated with Tasigna.
- if you develop chest pain or discomfort, numbness or weakness, problems with walking or with your speech, pain, discolouration or a cool feeling in a limb, **tell your doctor immediately** as this may be a sign of a cardiovascular event. Serious cardiovascular events including problems with the blood flow to the leg (peripheral arterial occlusive disease), ischaemic heart disease and problems with the blood supply to the brain (ischaemic cerebrovascular disease) have been reported in patients taking Tasigna. Your doctor should assess the level of fats (lipids) and sugar in your blood before initiating treatment with Tasigna and during treatment.
- if you develop swelling of the feet or hands, generalised swelling or rapid weight gain tell your doctor as these may be signs of severe fluid retention. Uncommon cases of severe fluid retention have been reported in patients treated with Tasigna.

If you are the parent of a child who is being treated with Tasigna, tell the doctor if any of the above

conditions apply to your child.

Children and adolescents

Tasigna is a treatment for children and adolescents with CML. There is no experience with the use of this medicine in children below 2 years of age. There is no experience with the use of Tasigna in newly diagnosed children below 10 years of age and limited experience in patients below 6 years of age who are no longer benefiting from previous treatment for CML. The long-term effects of treating children with Tasigna for long periods of time are not known.

Other medicines and Tasigna

Tasigna may interfere with some other medicines.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes in particular:

- anti-arrhythmics – used to treat irregular heart beat;
- chloroquine, halofantrine, clarithromycin, haloperidol, methadone, moxifloxacin - medicines that may have an unwanted effect on the function of the heart;
- ketoconazole, itraconazole, voriconazole, clarithromycin, telithromycin – used to treat infections;
- ritonavir – a medicine from the class “ antiproteases” used to treat HIV;
- carbamazepine, phenobarbital, phenytoin – used to treat epilepsy;
- rifampicin – used to treat tuberculosis;
- St. John’s Wort – a herbal product used to treat depression and other conditions (also known as *Hypericum perforatum*);
- midazolam – used to relieve anxiety before surgery;
- alfentanil and fentanyl – used to treat pain and as a sedative before or during surgery or medical procedures;
- cyclosporine, sirolimus and tacrolimus – medicines that suppress the “self-defence” ability of the body and fight infections and are commonly used to prevent the rejection of transplanted organs such as the liver, heart and kidney;
- dihydroergotamine and ergotamine – used to treat dementia;
- lovastatin, simvastatin – used to treat high level of fats in blood;
- warfarin – used to treat blood coagulation disorders (such as blood clots or thromboses);
- astemizole, terfenadine, cisapride, pimozone, quinidine, bepridil or ergot alkaloids (ergotamine, dihydroergotamine).

These medicines should be avoided during your treatment with Tasigna. If you are taking any of these, your doctor might prescribe other alternative medicines.

In addition, tell your doctor or pharmacist before taking Tasigna if you are taking any antacids, which are medicines against heartburn. These medicines need to be taken separately from Tasigna:

- H₂ blockers, which decrease the production of acid in the stomach. H₂ blockers should be taken approximately 10 hours before and approximately 2 hours after you take Tasigna;
- antacids such as those containing aluminium hydroxide, magnesium hydroxide and simethicone, which neutralise high acidity in the stomach. These antacids should be taken approximately 2 hours before or approximately 2 hours after you take Tasigna.

You should also tell your doctor **if you are already taking Tasigna** and you are prescribed a new medicine that you have not taken previously during Tasigna treatment.

Tasigna with food and drink

Do not take Tasigna with food. Food may enhance the absorption of Tasigna and therefore increase the amount of Tasigna in the blood, possibly to a harmful level. Do not drink grapefruit juice or eat grapefruit. It may increase the amount of Tasigna in the blood, possibly to a harmful level.

Pregnancy and breast-feeding

- **Tasigna is not recommended during pregnancy** unless clearly necessary. If you are pregnant

or think that you may be, tell your doctor who will discuss with you whether you can take this medicine during your pregnancy.

- **Women who might get pregnant** are advised to use highly effective contraception during treatment and for up to two weeks after ending treatment.
- **Breast-feeding is not recommended** during treatment with Tasigna. Tell your doctor if you are breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

If you experience side effects (such as dizziness or visual disorders) with a potential impact on the ability to safely drive or use any tools or machines after taking this medicine, you should refrain from these activities until the effect has disappeared.

Tasigna contains lactose

This medicine contains lactose (also known as milk sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Tasigna

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much Tasigna to take

Use in adults

- The recommended dose is 600 mg per day. This dose is achieved by taking two hard capsules of 150 mg twice a day.

Use in children and adolescents

- The dose given to your child will depend on your child's body weight and height. The doctor will calculate the correct dose to use and tell you how many capsules of Tasigna to give to your child. The total daily dose you give to your child must not exceed 800 mg.

Your doctor may prescribe a lower dose depending on how you respond to treatment.

Older people (age 65 years and over)

Tasigna can be used by people aged 65 years and over at the same dose as for other adults.

When to take Tasigna

Take the hard capsules:

- twice a day (approximately every 12 hours);
- at least 2 hours after any food;
- then wait 1 hour before eating again.

If you have questions about when to take this medicine, talk to your doctor or pharmacist. Taking Tasigna at the same time each day will help you remember when to take your hard capsules.

How to take Tasigna

- Swallow the hard capsules whole with water.
- Do not take any food together with the hard capsules.
- Do not open the hard capsules unless you are unable to swallow them. If so, you may sprinkle the content of each hard capsule in **one** teaspoon of apple sauce and take it immediately. Do not use more than one teaspoon of apple sauce for each hard capsule and do not use any food other than apple sauce.

How long to take Tasigna

Continue taking Tasigna every day for as long as your doctor tells you. This is a long-term treatment. Your doctor will regularly monitor your condition to check that the treatment is having the desired effect.

Your doctor may consider discontinuing your treatment with Tasigna based on specific criteria. If you have questions about how long to take Tasigna, talk to your doctor.

If you take more Tasigna than you should

If you have taken more Tasigna than you should have, or if someone else accidentally takes your hard capsules, contact a doctor or hospital for advice straight away. Show them the pack of hard capsules and this package leaflet. Medical treatment may be necessary.

If you forget to take Tasigna

If you miss a dose, take your next dose as scheduled. Do not take a double dose to make up for a forgotten hard capsule.

If you stop taking Tasigna

Do not stop taking this medicine unless your doctor tells you to do so. Stopping Tasigna without your doctor's recommendation places you at risk for worsening of your disease which could have life-threatening consequences. Be sure to discuss with your doctor, nurse, and/or pharmacist if you are considering stopping Tasigna.

If your doctor recommends that you discontinue treatment with Tasigna

Your doctor will regularly evaluate your treatment with a specific diagnostic test and decide whether you should continue to take this medicine. If you are told to discontinue Tasigna, your doctor will continue to carefully monitor your CML before, during and after you have discontinued Tasigna and may tell you to re-start Tasigna if your condition indicates that this is necessary.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the side effects are mild to moderate and will generally disappear after a few days to a few weeks of treatment.

Some side effects could be serious.

These side effects are common (may affect up to 1 in 10 people), uncommon (may affect up to 1 in 100 people) or have been reported with frequency not known (cannot be estimated from the available data).

- rapid weight gain, swelling of hands, ankles, feet or face (signs of water retention)
- chest pain, high blood pressure, irregular heart rhythm, blue discolouration of the lips, tongue or skin (signs of heart disorders)
- difficulty breathing, cough, wheezing with or without fever, swelling of the feet or legs (signs of lung disorders)
- fever, easy bruising, frequent infections (signs of blood disorders)
- blurred vision, loss of vision, blood in eye (signs of eye disorders)
- swelling and pain in one part of the body (signs of clotting within a vein)
- abdominal pain, nausea, constipation, swollen abdomen (signs of gastrointestinal disorders)
- severe upper abdominal pain (sign of pancreatitis)
- yellow skin and eyes, nausea, loss of appetite, dark-coloured urine (signs of liver disorders)
- rash, painful red lumps, pain in joints and muscles (signs of skin disorders)
- excessive thirst, high urine output, increased appetite with weight loss, tiredness (signs of high level of sugar in the blood)
- nausea, shortness of breath, irregular heartbeat, clouding of urine, tiredness and/or joint discomfort associated with abnormal results of blood tests (such as high levels of potassium,

- uric acid and phosphorous and low levels of calcium)
- pain, discomfort, weakness or cramping in the leg muscles, which may be due to decreased blood flow, ulcers on the legs or arms that heal slowly or not at all and noticeable changes in colour (blueness or paleness) or temperature (coolness) of the legs or arms, as these symptoms could be signs of artery blockage in the affected limb (leg or arm) and digits (toes and fingers)
- recurrence (reactivation) of hepatitis B infection when you have had hepatitis B in the past (a liver infection).

If you get any of these, **tell your doctor straight away.**

Some side effects are very common (may affect more than 1 in 10 people)

- headache
- tiredness
- muscle pain
- itching, rash
- nausea
- hair loss
- high blood level of bilirubin (liver function)
- high blood level of lipase (pancreas function)
- musculoskeletal pain, muscle pain, pain in extremity, pain in joints, bone pain and spinal pain upon discontinuing treatment with Tasigna

If any of these affects you severely, tell your doctor.

Some side effects are common (may affect up to 1 in 10 people)

- diarrhoea, vomiting, abdominal discomfort, stomach discomfort after meals, flatulence, swelling or bloating of the abdomen
- bone pain, pain in joints, muscle spasms, pain in extremity, back pain, pain or discomfort in the side of the body
- eye irritation, swelling, discharge, itching or redness, dry eye (signs of eye disorders)
- skin reddening, dry skin, acne, wart, decreased skin sensitivity, hives
- loss of appetite, disturbed sense of taste, weight increase
- insomnia, anxiety
- night sweats, excessive sweating, hot flushes
- dizziness, spinning sensation
- palpitations (sensation of rapid heart beat)

If any of these affects you severely, tell your doctor.

Some side effects are uncommon (may affect up to 1 in 100 people)

- skin pain
- swelling of eyelids
- nose bleed
- flu-like symptoms
- tingling or numbness
- visual disturbances
- feeling body temperature change (including feeling hot, feeling cold)
- thickened patches of red/silver skin (signs of psoriasis)
- sensitive teeth

If any of these affects you severely, tell your doctor.

The following other side effects have been reported with frequency not known (cannot be estimated from the available data):

- memory loss, disturbed or depressed mood, lack of energy, generally feeling unwell
- oral thrush, bacterial infection of the skin
- blister, skin cyst, oily skin, thinning of the skin, dark patches of skin, skin discolouration
- increased skin sensitivity
- bleeding, tender or enlarged gums
- runny or stuffy nose, sneezing
- dry mouth, sore throat, mouth sores

- trembling
- eye pain or redness, pain, itching of the eyelids
- painful and swollen joints (gout), muscle weakness
- unconsciousness
- difficulty and pain when passing urine, exaggerated sense of needing to urinate
- frequent urine output, abnormal urine colour
- haemorrhoids
- feeling of hardening in the breasts, heavy periods, nipple swelling
- appetite disorder, weight decreased
- severe headache often accompanied by nausea, vomiting and sensitivity to light
- heartburn
- breast enlargement in men
- symptoms of restless legs syndrome (an irresistible urge to move a part of the body, usually the leg, accompanied by uncomfortable sensations)

If any of these affects you severely, tell your doctor.

During Tasigna treatment, you may also have some abnormal blood test results such as low level of blood cells (white cells, red cells, platelets), high blood level of lipase or amylase (pancreas function), high blood level of bilirubin (liver function) or high blood level of creatinine (kidney function), low or high blood level of insulin (a hormone regulating blood sugar level), low or high level of sugar, or high level of fats in the blood.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Tasigna

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Store in the original package in order to protect from moisture.
- Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Tasigna contains

- The active substance is nilotinib. Each hard capsule contains 150 mg nilotinib (as hydrochloride monohydrate).
- The other ingredients are lactose monohydrate, crospovidone type A, poloxamer 188, colloidal anhydrous silica, magnesium stearate. The hard capsule shell is composed of gelatin, titanium dioxide (E171), red and yellow iron oxide (E172) and, shellac and black iron oxide (E172) for stamping of the imprint.

What Tasigna looks like and contents of the pack

Tasigna is supplied as hard capsules. The hard capsules are red. A black imprint is stamped on each hard capsule (“NVR/BCR”).

Tasigna is available in packs containing 28 or 40 hard capsules and in multipacks of 112 hard capsules

(comprising 4 cartons, each containing 28 hard capsules), 120 hard capsules (comprising 3 cartons, each containing 40 hard capsules) or 392 hard capsules (comprising 14 cartons, each containing 28 hard capsules).

Not all packs may be marketed in your country.

Marketing Authorisation Holder

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

Manufacturer

Novartis Pharma GmbH
Roonstraße 25
D-90429 Nuremberg
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Novartis Pharma N.V.
Tél/Tel: +32 2 246 16 11

България

Novartis Bulgaria EOOD
Тел.: +359 2 489 98 28

Česká republika

Novartis s.r.o.
Tel: +420 225 775 111

Danmark

Novartis Healthcare A/S
Tlf: +45 39 16 84 00

Deutschland

Novartis Pharma GmbH
Tel: +49 911 273 0

Eesti

Novartis Pharma Services Inc.
Tel: +372 66 30 810

Ελλάδα

Novartis (Hellas) A.E.B.E.
Τηλ: +30 210 281 17 12

España

Novartis Farmacéutica, S.A.
Tel: +34 93 306 42 00

France

Novartis Pharma S.A.S.
Tél: +33 1 55 47 66 00

Lietuva

Novartis Pharma Services Inc.
Tel: +370 5 269 16 50

Luxembourg/Luxemburg

Novartis Pharma N.V.
Tél/Tel: +32 2 246 16 11

Magyarország

Novartis Hungária Kft.
Tel.: +36 1 457 65 00

Malta

Novartis Pharma Services Inc.
Tel: +356 2122 2872

Nederland

Novartis Pharma B.V.
Tel: +31 26 37 82 555

Norge

Novartis Norge AS
Tlf: +47 23 05 20 00

Österreich

Novartis Pharma GmbH
Tel: +43 1 86 6570

Polska

Novartis Poland Sp. z o.o.
Tel.: +48 22 375 4888

Portugal

Novartis Farma - Produtos Farmacêuticos, S.A.
Tel: +351 21 000 8600

Hrvatska

Novartis Hrvatska d.o.o.
Tel. +385 1 6274 220

Ireland

Novartis Ireland Limited
Tel: +353 1 260 12 55

Ísland

Vistor hf.
Sími: +354 535 7000

Italia

Novartis Farma S.p.A.
Tel: +39 02 96 54 1

Κύπρος

Novartis Pharma Services Inc.
Τηλ: +357 22 690 690

Latvija

SIA "Novartis Baltics"
Tel: +371 67 887 070

România

Novartis Pharma Services Romania SRL
Tel: +40 21 31299 01

Slovenija

Novartis Pharma Services Inc.
Tel: +386 1 300 75 50

Slovenská republika

Novartis Slovakia s.r.o.
Tel: +421 2 5542 5439

Suomi/Finland

Novartis Finland Oy
Puh/Tel: +358 (0)10 6133 200

Sverige

Novartis Sverige AB
Tel: +46 8 732 32 00

United Kingdom

Novartis Pharmaceuticals UK Ltd.
Tel: +44 1276 698370

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu/>. There are also links to other websites about rare diseases and treatments.

Package leaflet: Information for the user

Tasigna 200 mg hard capsules nilotinib

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Tasigna is and what it is used for
2. What you need to know before you take Tasigna
3. How to take Tasigna
4. Possible side effects
5. How to store Tasigna
6. Contents of the pack and other information

1. What Tasigna is and what it is used for

What Tasigna is

Tasigna is a medicine containing an active substance called nilotinib.

What Tasigna is used for

Tasigna is used to treat a type of leukaemia called Philadelphia chromosome positive chronic myeloid leukaemia (Ph-positive CML). CML is a cancer of the blood which makes the body produce too many abnormal white blood cells.

Tasigna is used in adult and paediatric patients with newly diagnosed CML or in patients with CML who are no longer benefiting from previous treatment including imatinib. It is also used in adult and paediatric patients who experienced serious side effects with previous treatment and are not able to continue taking it.

How Tasigna works

In patients with CML, a change in DNA (genetic material) triggers a signal that tells the body to produce abnormal white blood cells. Tasigna blocks this signal, and thus stops the production of these cells.

Monitoring during Tasigna treatment

Regular tests, including blood tests, will be performed during treatment. These tests will monitor:

- the amount of blood cells (white blood cells, red blood cells and platelets) in the body to see how Tasigna is tolerated.
- pancreas and liver function in the body to see how Tasigna is tolerated.
- the electrolytes in the body (potassium, magnesium). These are important in the functioning of the heart.
- the level of sugar and fats in the blood.

The heart rate will also be checked using a machine that measures electrical activity of the heart (a test called an "ECG").

Your doctor will regularly evaluate your treatment and decide whether you should continue to take Tasigna. If you are told to discontinue this medicine, your doctor will continue to monitor your CML and may tell you to re-start Tasigna if your condition indicates that this is necessary.

If you have any questions about how Tasigna works or why it has been prescribed for you or your child, ask your doctor.

2. What you need to know before you take Tasigna

Follow all the doctor's instructions carefully. They may differ from the general information contained in this leaflet.

Do not take Tasigna

- if you are allergic to nilotinib or any of the other ingredients of this medicine (listed in section 6).

If you think you may be allergic, tell your doctor **before taking Tasigna**.

Warnings and precautions

Talk to your doctor or pharmacist before taking Tasigna:

- if you have suffered prior cardiovascular events such as a heart attack, chest pain (angina), problems with the blood supply to your brain (stroke) or problems with the blood flow to your leg (claudication) or if you have risk factors for cardiovascular disease such as high blood pressure (hypertension), diabetes or problems with the level of fats in your blood (lipid disorders).
- if you have a **heart disorder**, such as an abnormal electrical signal called "prolongation of the QT interval".
- if you are being **treated with medicines** that affect the heart beat (anti-arrhythmics) or the liver (see **Other medicines and Tasigna**).
- if you suffer from lack of potassium or magnesium.
- if you have a liver or pancreas disorder.
- if you have symptoms such as easy bruising, feeling tired or short of breath or have experienced repeated infections.
- if you have had a surgical procedure involving the removal of the entire stomach (total gastrectomy).
- if you have ever had or might now have a hepatitis B infection. This is because Tasigna could cause hepatitis B to become active again, which can be fatal in some cases. Patients will be carefully checked by their doctor for signs of this infection before treatment is started.

If any of these apply to you or your child, tell your doctor.

During treatment with Tasigna

- if you faint (loss of consciousness) or have an irregular heart beat while taking this medicine, **tell your doctor immediately** as this may be a sign of a serious heart condition. Prolongation of the QT interval or an irregular heart beat may lead to sudden death. Uncommon cases of sudden death have been reported in patients taking Tasigna.
- if you have sudden heart palpitations, severe muscle weakness or paralysis, seizures or sudden changes in your thinking or level of alertness, **tell your doctor immediately** as this may be a sign of a fast breakdown of cancer cells called tumour lysis syndrome. Rare cases of tumour lysis syndrome have been reported in patients treated with Tasigna.
- if you develop chest pain or discomfort, numbness or weakness, problems with walking or with your speech, pain, discolouration or a cool feeling in a limb, **tell your doctor immediately** as this may be a sign of a cardiovascular event. Serious cardiovascular events including problems with the blood flow to the leg (peripheral arterial occlusive disease), ischaemic heart disease and problems with the blood supply to the brain (ischaemic cerebrovascular disease) have been reported in patients taking Tasigna. Your doctor should assess the level of fats (lipids) and sugar in your blood before initiating treatment with Tasigna and during treatment.
- if you develop swelling of the feet or hands, generalised swelling or rapid weight gain tell your

doctor as these may be signs of severe fluid retention. Uncommon cases of severe fluid retention have been reported in patients treated with Tasigna.

If you are the parent of a child who is being treated with Tasigna, tell the doctor if any of the above conditions apply to your child.

Children and adolescents

Tasigna is a treatment for children and adolescents with CML. There is no experience with the use of this medicine in children below 2 years of age. There is no experience with the use of Tasigna in newly diagnosed children below 10 years of age and limited experience in patients below 6 years of age who are no longer benefiting from previous treatment for CML. The long-term effects of treating children with Tasigna for long periods of time are not known.

Other medicines and Tasigna

Tasigna may interfere with some other medicines.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes in particular:

- anti-arrhythmics – used to treat irregular heart beat;
- chloroquine, halofantrine, clarithromycin, haloperidol, methadone, moxifloxacin - medicines that may have an unwanted effect on the function of the heart;
- ketoconazole, itraconazole, voriconazole, clarithromycin, telithromycin – used to treat infections;
- ritonavir – a medicine from the class “ antiproteases” used to treat HIV;
- carbamazepine, phenobarbital, phenytoin – used to treat epilepsy;
- rifampicin – used to treat tuberculosis;
- St. John’s Wort – a herbal product used to treat depression and other conditions (also known as *Hypericum perforatum*);
- midazolam – used to relieve anxiety before surgery;
- alfentanil and fentanyl – used to treat pain and as a sedative before or during surgery or medical procedures;
- cyclosporine, sirolimus and tacrolimus – medicines that suppress the “self-defence” ability of the body and fight infections and are commonly used to prevent the rejection of transplanted organs such as the liver, heart and kidney;
- dihydroergotamine and ergotamine – used to treat dementia;
- lovastatin, simvastatin – used to treat high level of fats in blood;
- warfarin – used to treat blood coagulation disorders (such as blood clots or thromboses);
- astemizole, terfenadine, cisapride, pimozone, quinidine, bepridil or ergot alkaloids (ergotamine, dihydroergotamine).

These medicines should be avoided during your treatment with Tasigna. If you are taking any of these, your doctor might prescribe other alternative medicines.

In addition, tell your doctor or pharmacist before taking Tasigna if you are taking any antacids, which are medicines against heartburn. These medicines need to be taken separately from Tasigna:

- H2 blockers, which decrease the production of acid in the stomach. H2 blockers should be taken approximately 10 hours before and approximately 2 hours after you take Tasigna;
- antacids such as those containing aluminium hydroxide, magnesium hydroxide and simethicone, which neutralise high acidity in the stomach. These antacids should be taken approximately 2 hours before or approximately 2 hours after you take Tasigna.

You should also tell your doctor **if you are already taking Tasigna** and you are prescribed a new medicine that you have not taken previously during Tasigna treatment.

Tasigna with food and drink

Do not take Tasigna with food. Food may enhance the absorption of Tasigna and therefore increase the amount of Tasigna in the blood, possibly to a harmful level. Do not drink grapefruit juice or eat grapefruit. It may increase the amount of Tasigna in the blood, possibly to a harmful level.

Pregnancy and breast-feeding

- **Tasigna is not recommended during pregnancy** unless clearly necessary. If you are pregnant or think that you may be, tell your doctor who will discuss with you whether you can take this medicine during your pregnancy.
- **Women who might get pregnant** are advised to use highly effective contraception during treatment and for up to two weeks after ending treatment.
- **Breast-feeding is not recommended** during treatment with Tasigna. Tell your doctor if you are breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

If you experience side effects (such as dizziness or visual disorders) with a potential impact on the ability to safely drive or use any tools or machines after taking this medicine, you should refrain from these activities until the effect has disappeared.

Tasigna contains lactose

This medicine contains lactose (also known as milk sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Tasigna

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much Tasigna to take

Use in adults

- The recommended dose is 800 mg per day. This dose is achieved by taking two hard capsules of 200 mg twice a day.

Use in children and adolescents

- The dose given to your child will depend on your child's body weight and height. The doctor will calculate the correct dose to use and tell you how many capsules of Tasigna to give to your child. The total daily dose you give to your child must not exceed 800 mg.

Your doctor may prescribe a lower dose depending on how you respond to treatment.

Older people (age 65 years and over)

Tasigna can be used by people aged 65 years and over at the same dose as for other adults.

When to take Tasigna

Take the hard capsules:

- twice a day (approximately every 12 hours);
- at least 2 hours after any food;
- then wait 1 hour before eating again.

If you have questions about when to take this medicine, talk to your doctor or pharmacist. Taking Tasigna at the same time each day will help you remember when to take your hard capsules.

How to take Tasigna

- Swallow the hard capsules whole with water.
- Do not take any food together with the hard capsules.
- Do not open the hard capsules unless you are unable to swallow them. If so, you may sprinkle the content of each hard capsule in **one** teaspoon of apple sauce and take it immediately. Do not use more than one teaspoon of apple sauce for each hard capsule and do not use any food other

than apple sauce.

How long to take Tasigna

Continue taking Tasigna every day for as long as your doctor tells you. This is a long-term treatment. Your doctor will regularly monitor your condition to check that the treatment is having the desired effect.

Your doctor may consider discontinuing your treatment with Tasigna based on specific criteria.

If you have questions about how long to take Tasigna, talk to your doctor.

If you take more Tasigna than you should

If you have taken more Tasigna than you should have, or if someone else accidentally takes your hard capsules, contact a doctor or hospital for advice straight away. Show them the pack of hard capsules and this package leaflet. Medical treatment may be necessary.

If you forget to take Tasigna

If you miss a dose, take your next dose as scheduled. Do not take a double dose to make up for a forgotten hard capsule.

If you stop taking Tasigna

Do not stop taking this medicine unless your doctor tells you to do so. Stopping Tasigna without your doctor's recommendation places you at risk for worsening of your disease which could have life-threatening consequences. Be sure to discuss with your doctor, nurse, and/or pharmacist if you are considering stopping Tasigna.

If your doctor recommends that you discontinue treatment with Tasigna

Your doctor will regularly evaluate your treatment with a specific diagnostic test and decide whether you should continue to take this medicine. If you are told to discontinue Tasigna, your doctor will continue to carefully monitor your CML before, during and after you have discontinued Tasigna and may tell you to re-start Tasigna if your condition indicates that this is necessary.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the side effects are mild to moderate and will generally disappear after a few days to a few weeks of treatment.

Some side effects could be serious.

These side effects are common (may affect up to 1 in 10 people), uncommon (may affect up to 1 in 100 people) or have been reported with frequency not known (cannot be estimated from the available data).

- rapid weight gain, swelling of hands, ankles, feet or face (signs of water retention)
- chest pain, high blood pressure, irregular heart rhythm, blue discolouration of the lips, tongue or skin (signs of heart disorders)
- difficulty breathing, cough, wheezing with or without fever, swelling of the feet or legs (signs of lung disorders)
- fever, easy bruising, frequent infections (signs of blood disorders)
- weakness or paralysis of the limbs or face, difficulty speaking, severe headache, seeing, feeling or hearing things that are not there (signs of nervous system disorders)
- thirst, dry skin, irritability, dark urine, decreased urine output (signs of kidney disorders)
- blurred vision, loss of vision, blood in eye (signs of eye disorders)
- swelling and pain in one part of the body (signs of clotting within a vein)
- abdominal pain, nausea, vomiting of blood, black stools, constipation, swollen abdomen (signs of gastrointestinal disorders)
- severe upper abdominal pain (sign of pancreatitis)

- yellow skin and eyes, nausea, loss of appetite, dark-coloured urine (signs of liver disorders)
- rash, painful red lumps, pain in joints and muscles (signs of skin disorders)
- excessive thirst, high urine output, increased appetite with weight loss, tiredness (signs of high level of sugar in the blood)
- fast heartbeat, bulging eyes, weight loss, swelling at the front of the neck (signs of overactive thyroid gland)
- nausea, shortness of breath, irregular heartbeat, clouding of urine, tiredness and/or joint discomfort associated with abnormal results of blood tests (such as high levels of potassium, uric acid and phosphorous and low levels of calcium)
- pain, discomfort, weakness or cramping in the leg muscles, which may be due to decreased blood flow, ulcers on the legs or arms that heal slowly or not at all and noticeable changes in colour (blueness or paleness) or temperature (coolness) of the legs or arms, as these symptoms could be signs of artery blockage in the affected limb (leg or arm) and digits (toes and fingers)
- recurrence (reactivation) of hepatitis B infection when you have had hepatitis B in the past (a liver infection).

If you get any of these, **tell your doctor straight away**.

Some side effects are very common (may affect more than 1 in 10 people)

- diarrhoea
- headache
- tiredness
- muscle pain
- itching, rash
- nausea
- vomiting
- hair loss
- high blood level of bilirubin (liver function)
- high blood level of lipase (pancreas function)
- musculoskeletal pain, muscle pain, pain in extremity, pain in joints, bone pain and spinal pain upon discontinuing treatment with Tasigna

If any of these affects you severely, tell your doctor.

Some side effects are common (may affect up to 1 in 10 people)

- abdominal discomfort, stomach discomfort after meals, flatulence, swelling or bloating of the abdomen
- bone pain, pain in joints, muscle spasms
- pain including back pain, neck pain and pain in extremity, pain or discomfort in the side of the body
- eye irritation, swelling, discharge, itching or redness, dry eye (signs of eye disorders)
- skin reddening, dry skin, acne, wart, decreased skin sensitivity, hives
- loss of appetite, disturbed sense of taste, weight decrease or increase
- insomnia, depression, anxiety
- night sweats, excessive sweating, hot flushes
- dizziness, generally feeling unwell, spinning sensation
- tingling or numbness
- voice disorder
- nose bleed
- frequent urine output
- palpitations (sensation of rapid heart beat)

If any of these affects you severely, tell your doctor.

Some side effects are uncommon (may affect up to 1 in 100 people)

- increased skin sensitivity, skin pain
- swelling of the eyelids
- dry mouth, sore throat, mouth sores
- heartburn
- breast pain

- increased appetite
- attention disorder
- difficulty and pain when urinating, exaggerated sense of needing to urinate
- inability to achieve or maintain an erection
- breast enlargement in men
- flu-like symptoms, muscle weakness
- trembling
- decreased sharpness of vision
- severe headache often accompanied by nausea, vomiting and sensitivity to light
- visual disturbances
- oral or vaginal thrush
- muscle and joint stiffness
- unconsciousness
- weight gain
- feeling body temperature change (including feeling hot, feeling cold)
- thickened patches of red/silver skin (signs of psoriasis)
- sensitive teeth

If any of these affects you severely, tell your doctor.

The following other side effects have been reported with frequency not known (cannot be estimated from the available data):

- confusion, disorientation, memory loss, disturbed mood, lack of energy
- bacterial infection of the skin
- blister, skin cyst, oily skin, thinning of the skin, dark patches of skin, skin discolouration
- bleeding, tender or enlarged gums
- runny or stuffy nose, sneezing
- reddening and/or swelling and possibly peeling on the palms and soles (so called hand-foot syndrome)
- increased sensitivity of the eyes or the skin to light
- eye pain or redness, pain, itching of the eyelids
- difficulty hearing, ear pain, noises (ringing) in the ears
- painful and swollen joints (gout)
- blood in urine, abnormal urine colour, urinary incontinence
- haemorrhoids
- feeling of hardening in the breasts, heavy periods, nipple swelling
- symptoms of restless legs syndrome (an irresistible urge to move a part of the body, usually the leg, accompanied by uncomfortable sensations)

If any of these affects you severely, tell your doctor.

During Tasigna treatment, you may also have some abnormal blood test results such as low level of blood cells (white cells, red cells, platelets), high blood level of lipase or amylase (pancreas function), high blood level of bilirubin (liver function) or high blood level of creatinine (kidney function), low or high blood level of insulin (a hormone regulating blood sugar level), low or high level of sugar, or high level of fats in the blood.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Tasigna

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month.

- Do not store above 30°C.
- Store in the original package in order to protect from moisture.
- Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Tasigna contains

- The active substance is nilotinib. Each hard capsule contains 200 mg nilotinib (as hydrochloride monohydrate).
- The other ingredients are lactose monohydrate, crospovidone type A, poloxamer 188, colloidal anhydrous silica, magnesium stearate. The hard capsule shell is composed of gelatin, titanium dioxide (E171), yellow iron oxide (E172) and, shellac (E904) and red iron oxide (E172) for stamping of the imprint.

What Tasigna looks like and contents of the pack

Tasigna is supplied as hard capsules. The hard capsules are light yellow. A red imprint is stamped on each hard capsule (“NVR/TKI”).

Tasigna is available in a wallet containing 28 hard capsules and in a carton containing 28 or 40 hard capsules.

Tasigna is also available in multipacks of:

- 112 (4 wallets of 28) hard capsules.
- 112 (4 packs of 28) hard capsules.
- 120 (3 packs of 40) hard capsules.
- 392 (14 packs of 28) hard capsules.

Not all packs may be marketed in your country.

Marketing Authorisation Holder

Novartis Europharm Limited
Frimley Business Park
Camberley GU16 7SR
United Kingdom

Manufacturer

Novartis Pharma GmbH
Roonstraße 25
D-90429 Nuremberg
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Novartis Pharma N.V.
Tél/Tel: +32 2 246 16 11

Lietuva

Novartis Pharma Services Inc.
Tel: +370 5 269 16 50

България

Novartis Bulgaria EOOD
Тел.: +359 2 489 98 28

Luxembourg/Luxemburg

Novartis Pharma N.V.
Tél/Tel: +32 2 246 16 11

Česká republika

Novartis s.r.o.
Tel: +420 225 775 111

Danmark

Novartis Healthcare A/S
Tlf: +45 39 16 84 00

Deutschland

Novartis Pharma GmbH
Tel: +49 911 273 0

Eesti

Novartis Pharma Services Inc.
Tel: +372 66 30 810

Ελλάδα

Novartis (Hellas) A.E.B.E.
Τηλ: +30 210 281 17 12

España

Novartis Farmacéutica, S.A.
Tel: +34 93 306 42 00

France

Novartis Pharma S.A.S.
Tél: +33 1 55 47 66 00

Hrvatska

Novartis Hrvatska d.o.o.
Tel. +385 1 6274 220

Ireland

Novartis Ireland Limited
Tel: +353 1 260 12 55

Ísland

Vistor hf.
Sími: +354 535 7000

Italia

Novartis Farma S.p.A.
Tel: +39 02 96 54 1

Κύπρος

Novartis Pharma Services Inc.
Τηλ: +357 22 690 690

Latvija

SIA "Novartis Baltics"
Tel: +371 67 887 070

Magyarország

Novartis Hungária Kft.
Tel.: +36 1 457 65 00

Malta

Novartis Pharma Services Inc.
Tel: +356 2122 2872

Nederland

Novartis Pharma B.V.
Tel: +31 26 37 82 555

Norge

Novartis Norge AS
Tlf: +47 23 05 20 00

Österreich

Novartis Pharma GmbH
Tel: +43 1 86 6570

Polska

Novartis Poland Sp. z o.o.
Tel.: +48 22 375 4888

Portugal

Novartis Farma - Produtos Farmacêuticos, S.A.
Tel: +351 21 000 8600

România

Novartis Pharma Services Romania SRL
Tel: +40 21 31299 01

Slovenija

Novartis Pharma Services Inc.
Tel: +386 1 300 75 50

Slovenská republika

Novartis Slovakia s.r.o.
Tel: +421 2 5542 5439

Suomi/Finland

Novartis Finland Oy
Puh/Tel: +358 (0)10 6133 200

Sverige

Novartis Sverige AB
Tel: +46 8 732 32 00

United Kingdom

Novartis Pharmaceuticals UK Ltd.
Tel: +44 1276 698370

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu/>. There are also links to other websites about rare diseases and treatments.